

DBA/1

DBA/10laHsd

Developed in 1909 by CC Little from mice used in color experiments and this strain is the oldest of all inbred strains of mice. The DBA is the oldest of all inbred strains of mice.

In 1929-30, crosses were made between sub-lines, and several new lines established; two of these were called 12 (now DBA/1) and 212 (now DBA/2). In 1955, from the Jackson Laboratory, to Laboratory Animals Centre, Carshalton, UK.

In 1979, from Laboratory Animals Centre to OLAC (now Inotiv). Harlan became Envigo in 2015, then Envigo was acquired by Inotiv in 2021.

RESEARCH APPLICATIONS

Coat color, behavior, immunology, type II collagen-induced arthritis, etc.

CHARACTERISTICS

Anatomy

Low brain weight (Storer, 1967). High erythrocyte count, low mean corpuscular volume (Russell *et al*, 1951). Large number of Peyer's patches (Hummel *et al*, 1966).

Behavior

High food drive and high open-field activity (Thompson, 1953). Low open-field activity (Bruell, 1964). Good performance in food-seeking task (Henderson, 1970). Low preference for sweet tasting substances (saccharin, sucrose, dulcin and acesulfame, averaged) (Lush 1988). Bad performance in shuttle box test compared to DBA/2 (Yanovsky *et al*, 1995).

Drugs

Resistant to skin ulceration by DMBA (Thomas *et al*, 1973). Resistant to induction of subcutaneous tumors by 3-methylcholanthrene (Kouri *et al*, 1973; Whitmire *et al*, 1971). Sensitive to X-irradiation (Roderick, 1963). Males have a long sleeping time under hexobarbital (Lovell, 1976), long sleeping time under pentobarbitone anesthetic (Lovell, 1986). Insensitive (eosinophil response) to cortisone acetate (Wragg and Speirs, 1952). Sensitive to teratogenic effect (cleft palate) by cortisone acetate (Kalter 1981). Sensitive to seizures induced by nicotine (Marks *et al*, 1989). Clonidine induces a strong aggressive behavioral response (Nikulina and Klimek, 1993).

Genetics

Coat color genes

- *a, b, C, d* : non-agouti, dilute brown.

Histocompatibility

- *H-1^a, H-2^a, H-3^b, Thy-1^b*.

Biochemical markers

- *Apoa-1^b, Car-2^a, Es-1^b, Es-2^b, Es-3^c, Gpd-1^a, Gpi-1^a, Hbb^d, Idh-1^b, Ldr-1^a, Mod-1^a, Pep-3^b, Pgm-1^b, Trf^b*.

Although the DBA/1 and DBA/2 are substrains of the DBA there are differences between these strains, probably due to a substantial residual heterozygosity following the crosses between the substrains. DBA/1 and DBA/2 differ at least at the following loci: *Car-2, Ce-2, Hc, H-2, If-1, Lsh, Tla, and Qa-3*. With such large differences, they should probably be regarded as different strains rather than substrains of the same strain.

Immunology

Low lymphocyte phytohemagglutinin response (Heiniger *et al*, 1975). Poor immune response to ovomucoid, but good response to ovalbumin (Vaz *et al*, 1971). Good primary immune response to bovine serum albumin (James and Milne, 1972). Good primary immune response to sheep erythrocytes (Ghaffar and James, 1973). Non-discriminator between 'H' and 'L' sheep erythrocytes (McCarthy and Dutton, 1975). Poor immune response to (Pro-Gly-Pro)n (Fuchs *et al*, 1974). High susceptibility to IgG1-mediated but low susceptibility to IgE-mediated passive cutaneous anaphylaxis (De Souza *et al*, 1974). Good immune response to *Salmonella strasbourg* lipopolysaccharide (Di Pauli, 1972). Erythrocytes have a high agglutinability (Rubinstein *et al*, 1974). The DBA/1 was found to be useful on type II collagen-induced arthritis (Courtenay *et al*, 1980; Watson *et al*, 1985). Oral administration of *Lactobacillus casei* has a suppressive effect on type II collagen-induced arthritis in DBA/1 mice (Kato,





1998). *Staphylococcal enterotoxin B* induces arthritis in female DBA/1 mice but fails to induce activation of type II collagen-reactive lymphocytes (Omata *et al*, 1997). Study of the production of different cytokines during the development of collagen-induced arthritis (Mussener *et al*, 1997).

Infection

Susceptible to *Mycoplasma fermentans* (Gabridge *et al*, 1972). Resistant to *Plasmodium berghei* infection (Most *et al*, 1966). High mortality in a natural epizootic of ectromelia (Briody, 1966). Rapid immunological expulsion of *Trichinella spiralis* worms (Wakelin and Donachie 1980). Susceptible to *Leishmania* infection (Bradley, 1977). Resistant to *Cryptococcus neoformans* due to *Hc1* allele (Rhodes *et al*, 1980). Susceptible to the development of chronic Chagas' cardiomyopathy in postacute *Trypanosoma cruzi* infection (Rowland *et al*, 1992). Infection with larval *Echinococcus multilocularis* by transportal injection of hyatid homogenate results in a multivesiculation form of hyatid development. Protoscoleces are well developed (Nakaya *et al*, 1997). Sensitive to WA1 (Babesia-like piroplasm) (Moro *et al*, 1998). Sensitive to cytomegalovirus (Price *et al*, 1990).

Life-span and spontaneous disease

Median life-span 14.4 months in DBA/1 males and 25.0 months in DBA/1 females (Storer, 1966). Median life-span 16.2 months in DBA/1 males and 24.0 months in DBA/1 females (Festing and Blackmore, 1971).

Primary lung tumors 3% in males, 1% in breeding females and zero in virgin females; lymphatic leukemia less than 1%. Mammary adenocarcinomas zero in males, 90% in breeding females and 61% in virgin females in unfostered substrain (Hoag, 1963).

A high proportion of the mammary tumors are of the acinar type (Tengbergen, 1970). Lung tumors 2-27% (Festing and Blackmore, 1971). Low gross tumor incidence in males (Storer, 1966). Mammary changes in female DBA/1 mice have been described by Orellana *et al* (2000).

Miscellaneous

DBA/1 mice are resistant to most of the DBA/2 tumors; lymphatic leukemia P1534 grows in 50% of DBA/1 mice, melanoma S91 grows in both strains (Staats, 1985). Recommended host for the following transplantable tumors:

anaplastic carcinoma dbrB, mammary adenocarcinomas CaDI and T1703, melanoma S91 and pleomorphic sarcoma S37 (which is not host-specific) (Kaliss, 1972). An embryonic stem cell line has been developed by Roach *et al* (1995). High incidence of spontaneous 'deviants' (possible mutations) (Schlager and Dickie, 1967). Characteristics of the DBA/2 strain have been described by Festing (1997) and Lyon *et al*, (1996).

Physiology and biochemistry

High serum ceruloplasmin levels (Meier and MacPike, 1968). High plasma cholinesterase activity in females (males not measured) (Angel *et al*, 1967). Low liver tyrosine aminotransferase in fasted mice (Blake, 1970). Low cell turnover as estimated by slow clearance of DNA-bound radioactivity (Heiniger *et al*, 1972). Low venous and arterial blood pH (Bernstein, 1966). A comparative study of the *in vitro* development of embryos (Suzuki *et al*, 1996)

Reproduction

Poor breeding performance, colony output 0.77 young/female/week, litter size 4.4 weaned (Festing, 1976).



REFERENCES

- Angel CR, Mahin DT, Farris RD, Woodward KT (1967) Heritability of plasma cholinesterase activity in inbred mouse strains. *Science* 156, 529-530.
- Bernstein SE (1966) Physiological characteristics. In: *Biology of the Laboratory Mouse*. 2nd ed. (Green EL, ed). New York: McGraw-Hill, pp. 337-350.
- Blake RL (1970) Regulation of liver tyrosine amino transferase activity in inbred strains and mutant mice. I. Strain variance in fasting enzyme levels. *Int. J. Biochem.* 1, 361-370.
- Bradley DJ (1977) Regulation of the leishmania populations within the host. II. Genetic control of acute susceptibility of mice to *L. donovani* infection. *Clin. Exp. Immunol.* 30, 130-140.
- Bridoy BA (1966) The natural history of mouse pox. *National Cancer Institute Monograph* 20, 105-116.
- Bruell JH (1964) Inheritance of behavioural and physiological characters of mice and the problem of heterosis. *Am. Zool.* 4, 125-138.
- Courtenay JS, Dallman MJ, Dayan AD, Martin A, Mosedale B (1980) Immunization against heterologous type II collagen induces arthritis in mice. *Nature* 283, 666-668.
- De Souza CM, Maia LCS, Vaz NM (1974) Susceptibility to cutaneous anaphylaxis in inbred strains of mice. *J. Immunol.* 112, 1369-1372.
- Di Pauli R (1972) Genetics of the immune response. I. Differences in the specificity of antibodies to lipopolysaccharides among different strains of mice. *J. Immunol.* 109, 394-400.
- Festing MFW (1976) Breeding performance of mouse colonies at the MRC Laboratory Animals Centre. Unpublished data.
- Festing MFW (1997) Inbred Strains of mice. *Mouse genome* 95, 519-686.
- Festing MFW, Blackmore DK (1971) Life span of specified-pathogen-free (MRC category 4) mice and rats. *Lab. Anim.* 5, 179-192.
- Fuchs S, Mozes E, Maoz A, Sela M (1974) Thymus independence of a collagen-like synthetic polypeptide and of collagen and the need for thymus and bone marrow-cell cooperation in the immune response to gelatin. *J. Exp. Med.* 139, 148-158.
- Gabridge MG, Abrams GD, Murphy WH (1972) Lethal toxicity of *Mycoplasma fermentans* in mice. *J. Infect. Dis.* 125, 153-160.
- Ghaffar A, James K (1973) The effect of antilymphocyte antibody on the humoral immune response in different strains of mice. *Immunol.* 24, 455-465.
- Heiniger HJ, Chen HW, Meier H, Taylor BA, Commerford LS (1972) Studies on the genetic control of cell proliferation. I. Clearance of DNA-bound radioactivity in 19 inbred strains and hybrid mice. *Life Sci.* 11, 87-98.
- Heiniger HJ, Taylor BA, Hards EJ, Meier H (1975) Heritability of the phytohemagglutinin responsiveness of lymphocytes and its relationship to leukemogenesis. *Cancer Res.* 35, 825-831.
- Henderson ND (1970) Genetic influences on the behaviour of mice can be obscured by laboratory rearing. *J. Comp. Physiol. Psychol.* 72, 505-511.
- Hoag WG (1963) Spontaneous cancer in mice. *Ann. NY Acad. Sci.* 108, 805-831.
- Hummel KP, Richardson FL, Fekete E (1966) *Anatomy*. In: *Biology of the Laboratory Mouse*. 2nd ed. (Green EL, ed). New York: McGraw-Hill, pp. 247-307.
- James K, Milne I (1972) The effect of anti-lymphocytic antibody on the humoral immune response in different strains of mice. I. The response to bovine serum albumin. *Immunol.* 23, 897-909.
- Kaliss N (1972) Transplanted tumors. *Jax Notes* 410, Jackson Laboratory, Bar Harbor, Maine.
- Kaiter H (1981) Dose response studies with genetically homogeneous lines of mice as a teratology testing and risk assessment procedure. *Teratology* 24, 79-86.
- Kato I (1998) Suppressive effects of the oral administration of *Lactobacillus casei* on type II collagen-induced arthritis in DBA/1 mice. *Life Sciences* 63, 635-644.
- Kouri RE, Salerno RA, Whitmire CE (1973) Relationships between arylhydrocarbon hydroxylase inducibility and sensitivity to chemically induced subcutaneous sarcomas in various strains of mice. *J. Natl. Cancer Inst.* 50, 363-368.
- Lovell DP (1976) Hexobarbital sleeping time in mice. Unpublished data.
- Lovell DP (1986) Variation in pentobarbitone sleeping time in mice. I. strain and sex differences. *Lab. Anim.* 20, 85-90.
- Lush IM (1988) The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet. Res.* 53, 95-99.
- Lyon MF, Rastan S, Brown SDM (1996) Genetic variants and strains of the laboratory mouse. 2 Volumes. Oxford, New York, Tokyo: Oxford University Press.
- Marks MJ, Stitzel JA, Collins AC (1989) Genetic influences on nicotine responses. *Pharmacol. Biochem. Behav.* 33, 667-678.
- McCarthy MM, Dutton RW (1975) The humoral response of mouse spleen cells to two types of sheep erythrocytes. *J. Immunol.* 115, 1316-1321.
- Meier H, MacPike AD (1968) Levels and heritability of serum ceruloplasmin activity in inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 128, 1185-1190.
- Moro MH, David,CS, Magera JM, Wettstein PJ, Barthold SW, Persing DH (1998) Differential Effects of Infection with a Babesia-Like Piroplasm, WAI, in Inbred Mice. *Infect. Immun.* 66, 492-498.
- Most H, Nussenzeig RS, Vanderberg J, Herman R, Yoeli M (1966) Susceptibility of genetically standardized (JAX) mouse strains to sporozoite and blood-induced *Plasmodium berghei* infections. *Mil. Med.* 131, 915-918.
- Mussener A, Litton MJ, Lindroos E, Klareskog L (1997) Cytokine production in synovial tissue of mice with collagen-induced arthritis (CIA). *Clin. Exp. Immunol.* 107, 485-493.
- Nakaya K, Nakao M, Ito A (1997) *Echinococcus multilocularis*: Mouse strain difference in hydatid development. *J. Helminthology* 71, 53-56.
- Nikulina EM, Klimek V (1993) Strain differences in clonidine-induced aggressiveness in mice and its interaction with the dopamine system. *Pharmacol. Biochem. Behav.* 44, 821-825.
- Omata S et al (1997) Staphylococcal enterotoxin B induces arthritis in female DBA/1 mice but fails to induce activation of type II collagen-reactive lymphocytes. *Cellular Immunology* 179, 138-145.
- Orellana JM, Garcia MJ, Diez JC (2000) Histological differences in mammary glands of female DBA/1 mice. In: *Proceedings of the International Joint Meeting XII ICLA General Assembly & 7th FELASA symposium*, Palma de Mallorca, Spain. Laboratory Animals Ltd, London, pp.114-116.
- Price P, Gibbons AE, Shellam GR (1990) H-2 class I loci determine sensitivity to MCMV in macrophages and fibroblasts. *Immunogenetics* 32, 20-26.
- Rhodes JC, Wicker LS, Urba WJ (1980) Genetic control of susceptibility to *Cryptococcus neoformans* in mice. *Infect Immun* 29, 494-499.
- Roach ML, Stock JL, Byrum R, Koller BH, McNeish JD (1995) A new embryonic stem cell line from DBA/1lacJ mice allows genetic modification in a murine model of human inflammation. *Exp. Cell Research* 221, 520-525.
- Roderick TH (1963) The response of twenty-seven inbred strains of mice to daily doses of whole-body X-irradiation. *Radiation Res.* 20, 631-639.
- Rowland EC, Lozykowski MG, McCormick TS (1992) Differential cardiac histopathology in inbred mouse strains chronically infected with *Trypanosoma cruzi*. *Journal of Parasitology* 78, 1059-1066.
- Rubinstein P, Liu N, Strenn EW, Decary F (1974) Electrophoretic mobility and agglutinability of red blood cells: a 'new' polymorphism in mice. *J. Exp. Med.* 139, 313-322.
- Russell ES, Neufeld EF, Higgins CT (1951) Comparison of normal blood picture of young adults from 18 inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 78, 761-766.
- Schlager G, Dickie MM (1967) Spontaneous mutations and mutation rates in the house mouse. *Genetics* 57, 319-330.
- Staats J (1985) Standardized nomenclature for inbred strains of mice: eighth listing. *Cancer Research* 45, 945-977.
- Storer JB (1966) Longevity and gross pathology at death in 22 inbred strains of mice. *J. Gerontol.* 21, 404-409.
- Storer JB (1967) Relation of lifespan to brain weight, body weight and metabolic rate among inbred mouse strains. *Exp. Gerontol.* 2, 173-182.
- Suzuki O, Asano T, Yamamoto Y, Takano K, Koura M (1996) Development in vitro of preimplantation embryos from 55 mouse strains. *Reproduction, Fertility and Development.* 8, 975-980.
- Tengbergen WJPR van E (1970) Morphological classification of mammary tumors in the mouse. *Path. Eur.* 5, 260-272.
- Thomas PE, Hutton JJ, Taylor BA (1973) Genetic relationship between aryl hydrocarbon hydroxylase inducibility and chemical carcinogen induced skin ulceration in mice. *Genetics* 74, 655-659.
- Thompson WR (1953) The inheritance of behaviour: behavioural differences in fifteen mouse strains. *Can. J. Psychol.* 7, 145-155.
- Vaz NM, Phillips-Quagliata JM, Levine BB, Vaz EM (1971) H-2 linked genetic control of immune responsiveness to ovalbumin and ovomucoid. *J. Exp. Med.* 134, 1335-1348.
- Wakelin D, Donachie AM (1980) Genetic control of immunity to parasites: adoptive transfer of immunity between inbred strains of mice characterized by rapid and slow immune expulsion of *Trichinella spiralis*. *Parasite Immunol.* 2, 249-260.
- Watson WC et al (1985) Genetic susceptibility to murine collagen II autoimmune arthritis. *J. Exp. Med.* 162, 1878-1891.
- Whitmire CE, Salerno RA, Rabstein LS, Heubner RJ, Turner HC (1971) RNA tumor-virus antigen expression in chemically induced tumors. Virus-genome specified common antigens detected by complement fixation in mouse tumours induced by 3-methylcholanthrene. *J. Natl. Cancer Inst.* 47, 1255-1265.
- Wragg LE, Speirs RS (1952) Strain and sex differences in response of inbred mice to adrenal cortical hormones. *Proc. Soc. Exp. Biol. Med.* 80, 680-684.
- Yanovsky Y, Brankack J, Haas HL (1995). Differences of CA3 bursting in DBA/1 and DBA/2 inbred mouse strains with divergent shuttle box performance. *Neuroscience* 64, 319-325.