

C3H

C3H/HeNHsd

Developed in 1920 by Strong from a cross of Bagg albino female and DBA male. Strain CBA also originated from this cross. Strain CBA was selected for a low mammary tumor incidence and C3H for a high incidence. In 1930, to Andervont, National Cancer Institute, Bethesda, MD, USA and in 1941, to Heston and Law at the same institute.

Derived from a nucleus colony obtained from the National Institutes of Health, Bethesda, Maryland.

RESEARCH APPLICATIONS

Behavior, aggression, mammary tumors, metabolism, immunology, infectious diseases, parasitology, etc.

CHARACTERISTICS

Strain C3H was selected for a high mammary tumor incidence. Unfostered sub-lines have a very high incidence of mammary tumors caused by a virus (Bittner virus), which is transmitted, from mother to offspring through the milk. Fostering of the young to a mammary tumor virus-free strain eliminates the virus and therefore all Germfree or SPF (Specified Pathogen Free) C3H mice will be free of this virus. Fostered C3H mice are widely used as a general-purpose strain.

Anatomy

Low total leukocyte count, low erythrocyte count, low hematocrit, and low hemoglobin (Russell *et al*, 1951). Small thymus/body weight ratio (Belyaev *et al*, 1970), small thymus/body weight ratio (depending on age) (Albert *et al*, 1965). Large pituitary (Sinha *et al*, 1975). Adrenal gland X zone large, with high incidence of vacuolization (Delost and Chirvan-Nia, 1958). Low percentage of mice has accessory spleens (Hummel *et al*, 1966). Many Peyer's patches (Hummel *et al*, 1966). Plasma volume low at 5.97 ml/100 g and red cell volume low at 4.64 ml/100g body weight in He substrain (Kano and Mizuma, 1974). High bone density of femur (Beamer *et al*, 1996).

Behavior

C3H mice are easy to handle. Low intra-strain aggression (Southwick and Clark, 1966). Long latency to emerge from home-cage, low rearing, long latency to cross barrier in open-field, low hole-in-the-wall entries and low Y-maze exploration (McClearn *et al*, 1970). Low open-field activity (Bruell, 1964). Low open-field defecation (Bruell, 1969). High food drive (Thompson, 1953), but poor performance in food-seeking task (Henderson, 1970). Short time of immobility in a forced swimming test (Nikulina *et al*, 1991). Low shock-avoidance learning (Bovet *et al*, 1966; Bovet *et al*, 1969). Good short-term but poor long-term memory in contrast with DBA/2 (Bovet *et al*, 1969). Good T-maze learning (Stasik, 1970). Poor water-escape

learning (Festing, 1973). Low radial-arm maze learning (Ammassari-Teule *et al*, 1993). High social grooming scores during aggressive encounters (Southwick and Clark, 1968). Carries the retinal degeneration gene and is capable of pattern discrimination up to 40 days, and brightness discrimination to at least 100 days (Nagy and Misanin, 1970).

Drugs

Susceptible to skin ulceration to DMBA (Thomas *et al*, 1973). Sensitive to the development of uterine tumors following treatment with DMBA at four weeks of age (Tsubura *et al*, 1993). Susceptible to induction of subcutaneous tumors by 3-methylcholanthrene, depending on substrain (Kouri *et al*, 1973). Susceptible to tumor induction by 3-methylcholanthrene in fostered and unfostered substrains (Whitmire *et al*, 1971; Whitmire and Salerno, 1972). Susceptible to induction of liver but resistant to pulmonary tumors by neonatally administered DMBA (Flaks, 1968). High susceptibility to tumor induction by 3,4-benzopyrene (Liebelt *et al*, 1970). High susceptibility to induction of mammary tumors by urethane (Bentvelzen *et al*, 1970). High incidence of gastric tumors after administration of methylcholanthrene by gavage (Akamatsu and Barton, 1974). Susceptible to fibrosarcoma induction by methylcholanthrene (Strong, 1952). Highly susceptible to the induction of hepatocellular tumors by various carcinogens, with the volume of hepatic





lesions being >100-fold greater than in more resistant strains. Susceptibility is linked to at least six chromosomal regions (Dragani *et al*, 1995). Phenobarbitone in the diet to give an intake of 85 mg/kg per day resulted in 70% of animals developing basophilic nodules by 91 weeks of age (contrast 4% in C57BL/6), but no increase in liver carcinomas (Evans *et al*, 1992). However, there was a two-fold greater level of DNA synthesis in C3H mice relative to C57BL/6 mice after partial hepatectomy, though partial hepatectomy is a tumor promoter in C57BL/6 but not in C3H mice (Bennett *et al*, 1995). Insensitive to histamine (Brown, 1965). Airways of C3H/HeJ hyporeactive to acetylcholine (Zhang *et al*, 1995). Resistant to teratogenic effect of acetazolamide (Green *et al*, 1973). Pentobarbital i.p. induces hepatic epoxide hydrase (Oesch *et al*, 1973). Sensitive to X-irradiation (Roderick, 1963). Long survival on Warfarin (Lush and Arnold, 1975). Sensitive to hyperbaric oxygen (Hill *et al*, 1968). Sensitive uterine response to estrogen's (Chai and Dickie, 1966). Short hexobarbital sleeping time (Vesell, 1968). Long survival in 90% oxygen and highly susceptible to pulmonary hyaline-membrane formation (Lieberman and Kellog, 1967). Resistant to the induction of pulmonary fibrosis by bleomycin (contrast C57BL/6) (Haston *et al*, 1996), and irradiation though the sensitivity of lung fibroblasts to irradiation in-vitro does not correlate with in-vivo sensitivity (Dileto and Travis, 1996). Sensitive to chloroform toxicity (Deringer *et al*, 1953). Susceptible to toxic effects of isoniazid (Taylor, 1976). High ED50 to behavioral effects of nicotine (Marks *et al*, 1989). Low self-selection of nicotine, which is inversely correlated with sensitivity to nicotine-induced seizures (Robinson

et al, 1996). Low bronchial reactivity to methacholine and serotonin (Konno *et al*, 1993). No increase in renal lipid peroxidation following treatment with nickel (Misra *et al*, 1991). Susceptible to biliary tract injury following oral dosing with 500 micrograms of the fungal toxin sporidesmin (Bhathal *et al*, 1990).

Low histamine release from peritoneal mast cells induced by compound 48/80, a calcium dependent histamine releaser (Toda *et al*, 1989). High histamine release from peritoneal mast cells induced by Ca²⁺ ionophore A23187 (contrast C57BL/6) (Toda *et al*, 1989). Cadmium highly hepatotoxic (Shaikh *et al*, 1993). Resistant to ozone-induced decreases of tracheal potential (Takahashi *et al*, 1995, Kleeberger *et al*, 1993). Susceptible to weight loss induced by cocaine, but this is attenuated by anisomycin (Shimosato *et al*, 1994). Estrogen does not induce an increase in VLDL and LDL-cholesterol (like BALB/c, contrast C57BL/6 and C57L) (Srivastava, 1995).

Genetics

Coat colour genes

- *A, B, C, D* : agouti (wild type).

Histocompatibility

- *H-2^k, Thy-1^b*.

Biochemical markers

- *Apoa-1^b, Car-2^b, Es-1^b, Es-2^b, Es-3^c, Gpd-1^b, Gpi-1^b, Hba^c, Hbb^d, Idh-1^a, Ldr-1^a, Mod-1^a, Mup-1^a, Pep-3^b, Pgm-1^b, Pgm-2^a, Trf^b*.

This strain carries the *Mus musculus musculus* Y-chromosome, while others have the *M. m. domesticus* type (Nishioka, 1987).

Immunology

Sensitive to amyloid induction but low level of spontaneous amyloid formation (Ram *et al*, 1969). Low lymphocyte phytohemagglutinin response (Heiniger *et al*, 1975). Good immune response to small doses of bovine gamma globulin (Levine and Vaz, 1970). Poor immune response to Cholera A and B antigens (Cerny *et al*, 1971). Good splenic PFC immune response to pneumococcal polysaccharide (Amsbaugh *et al*, 1972). Immune response of SJL mice to type-III pneumococcal polysaccharide declines by 42 weeks, in contrast to BALB/c and C3H (Smith, 1976). Females fail to reject male skin grafts after 100 days (contrast nine strains) (Gasser and Silvers, 1971). Poor immune response to ovomucoid and ovalbumin (Vaz *et al*, 1971). Poor primary immune response to bovine serum albumin (James and Milne, 1972). Good immune response to *Salmonella anatum*, *S. senftenberg* and *S. strasbourg* lipopolysaccharide (Di Pauli, 1972). Responder to synthetic polypeptide Glu⁵⁷, Lys³⁸, Ala⁵ (Pinchuk and Maurer, 1965). Good immune response to Vi antigen (Gaines *et al*, 1965). Precipitating and skin-sensitising antibodies have slow electrophoretic mobility (Fahey, 1965). High antibody affinity to HSA (Petty *et al*, 1972).

Erythrocytes have high agglutinability (Rubinstein *et al*, 1974). Low immune response to ferritin in He substrain (Young *et al*, 1976). Non-discriminator between 'H' and 'L' sheep erythrocytes (McCarthy and Dutton, 1975). High anti-DNP antibody concentration (Paul *et al*, 1970). Antibodies to lipoid A antigen do not cross-react with sheep red blood cells (contrast eight strains). Strain also resistant to toxic effect of *Salmonella*

lipopolysaccharide (Rank *et al*, 1969). Refractory to sensitising effects of HSF from *Bordetella pertussis* to histamine (contrast sixteen strains) (Bergman and Munoz, 1968). Good immune response to Pro-Gly-Pro-ovalbumin and (Pro-Gly-Pro)n (Fuchs *et al*, 1974). High susceptibility to IgE-mediated passive cutaneous anaphylaxis (De Souza *et al*, 1974). He substrain resistant to induction of anaphylactic shock by ovalbumin (Tanioka and Esaki, 1971). He and HeN substrains are susceptible to experimental autoimmune orchitis induced by two or three sc injections with viable syngeneic testicular germ cells without any adjuvants, but C3H/BiKi is resistant (Tokunaga *et al*, 1993). High immune response to ganglio-series gangliosides in C3H/HeN, but low response in C3H/HeJ (Kawashima *et al*, 1992). Anti-BPO IgE monoclonal antibody did not produce potent systemic sensitisation sufficient for provocation of lethal shock in most aged (six-ten months) mice (Harada *et al*, 1991). Carries a strain-specific allele at the alpha globin locus (Sato *et al*, 1996). High natural killer cell response to the immunostimulant 7-allyl-8-oxoguanosine (Pope *et al*, 1994).

Infection

Resistant to infection by *Salmonella typhimurium* strain C5 (Plant and Glynn, 1974). Susceptible to *Mycoplasma fermentans* (Gabridge *et al*, 1972). Experimental *Mycoplasma pulmonis* infection results in acute pneumonia with severe hemorrhage, edema and often death (Faulkner *et al*, 1995). Susceptible to mammary tumor virus, which is carried in an active form in unfostered substrains (Murray and Little, 1967). Susceptible to oncogenic effect of polyoma virus given at birth (Law, 1966). Susceptible to measles virus induced encephalitis, which correlates with a high cytotoxic T-lymphocyte response (like C57BL/6, contrast BALB/c) (Niewiesk *et al*, 1993). Susceptible to *Mycobacterium marinum* (Shepard and Habas, 1967). Susceptible to infection by *Mycobacterium marinum* (Yamamoto *et al*, 1991). Susceptible to infection by *Entamoeba histolytica* (Neal and Harris, 1975). Resistant to mouse hepatitis virus (Bang and Warwick, 1960). 100% transmission of murine leukemia virus (Scripps) through three successive generations (Jenson *et al*, 1976). Highly susceptible to measles virus (Rager-

Zisman *et al*, 1976). Highly susceptible to tumor induction by polyoma virus (Freund *et al*, 1992). Following administration of murine cytomegalovirus, C3H mice exhibited minimal carditis after neonatal or adult infection. However neonatal infection appears to accelerate age-related cardiopathy, which is severe in retired breeders of this strain. (contrast BALB/c and C57BL/10) (Price *et al*, 1991). Highly susceptible to Lyme borreliosis (*Borrelia burgdorferi*) when inoculated at three weeks of age and as adults. Mice inoculated at age three weeks also developed polyarthritis, but severity was reduced when inoculated as adults. Carditis was also common (Barthold *et al*, 1990), and mice were susceptible to the development of arthritis (contrast BALB/c) (Matyniak and Reiner, 1995). Resistant to intra-vaginally inoculated *Neisseria gonorrhoea* (Johnson *et al*, 1989). Resistant to infection with *Ehrlichia risticii* (Williams and Timoney, 1994). Susceptible to infection by *Helicobacter felis* with moderate to severe chronic active gastritis in the body of the stomach, which increased over time (Sakagami *et al*, 1996).

Life-span and spontaneous disease

Almost 100% of mammary tumors in females of unfostered substrains (Heston, 1963). Mammary adenocarcinomas in unfostered substrains less than 1% in males, 95% in breeding and 88% in virgin females. Lymphatic leukemia zero incidence (Hoag, 1963). Mammary tumors 40% at 18.8 months in C3HfC57BL, but 99% at 7.2 months in unfostered C3H (Heston and Vlahakis, 1971). Mammary tumors 37% at two years in fostered substrain (Bentvelzen *et al*, 1970). Median latent period to develop mammary tumors in unfostered substrains ranged from 276 to 566 days, depending on breeding status and environmental stress (Riley, 1975). A high proportion of the mammary tumors are of the acinar type (Tengbergen, 1970). Incidence of mammary tumors reduced by bromocriptine and interferon Stravoravdi *et al*, 1991). Hepatomas 72-91% in males at 14 months, 59% in virgin females, 30-38% in breeding females (Heston, 1963). Hepatomas have eosinophilic cytoplasmic inclusion bodies (Liebelt *et al*, 1971). Good model of genetic predisposition to hepatocellular tumors, susceptibility being associated with six

chromosomal regions (Dragani *et al*, 1995). Point mutations in H-ras do not generally play a major or initiating role in spontaneous hepatocarcinogenesis in this strain (Enomoto *et al*, 1993). Lung adenomas 2-10% in fostered A substrain, leukemia 6-30% (Muhlbock and Tengbergen, 1971). Occasional Harderian gland tumors (Heston, 1963). Rare "lipomatous" hamartomas or choristomas have been noted (Adkison and Sundberg, 1991). Life-span in SPF fostered conditions intermediate in both sexes (590 days in males, 676 days in females). Liver tumors 9-23%, lung tumors 2-10% and mammary tumors 21-36%. Heart defects 13-26% and cystic ovaries 13-26% (Festing and Blackmore, 1971). The relationship of genotype, sex, body weight, and growth parameters to lifespan in inbred and hybrid mice is described by Ingram *et al* (1982). The influence of MHC on aging is described by Smith and Walford (1977). Mean survival 98 ± 3.0 weeks. The influence of the H-2 and H-1 histocompatibility upon life-span and cancer incidence is described by Smith and Walford (1978). Tail lesions similar in appearance to bite wounds were found in grouped C3H/HeJ by Les (1972). Develop dystrophic cardiac calcification, which may be related to disturbed myocyte calcium metabolism (Brunnert, 1997). Can be made obese by a suitable diet (Fenton and Dowling, 1953).

Resistant to the development of aortic cartilaginous metaplasia (contrast C57BL/6) (Qiao *et al*, 1995). Resistant to diet-induced aortic fatty streak lesions which correlates with a high level of paroxinase mRNA (contrast C57BL/6) (Shih *et al*, 1996).

Miscellaneous

Recommended host for the following transplantable tumors: lymphosarcoma 6C3HED and mammary adenocarcinomas C3HBA and H2712 (Kaliss, 1972). Recommended host for sarcoma BP8 used as a model for screening potential anticancer drugs (EORTC Screening Group, 1972). High mortality after neonatal thymectomy (Law, 1966). High rate of spontaneous mutations and total deviants (Schlager and Dickie, 1967). Characteristics of the C3H strain have been described by Festing (1997) and Lyon *et al*, (1996).



Physiology and biochemistry

Low serum calcium in Fg substrain but He substrain has high level at 4 months (Barrett *et al*, 1975). High serum cholesterol (Bruell *et al*, 1962). High plasma cholesterol and triglycerides. High erythrocyte catalase (Hoffman and Rechcigl, 1971). Low serum haptoglobin level (Peacock *et al*, 1967). Low peripheral nerve conduction velocity (Hegmann, 1972). Low percentage of time spent sleeping with low percentage of slow-wave sleep and small diurnal variation (Valatx and Bugat, 1974). Low metabolic rate (Pennycuik, 1967). High liver tyrosine aminotransferase level in fasted mice (Blake, 1970). Low adrenal corticosteroid production (Nandi *et al*, 1967). High peptidyl proline hydroxylase activity in tumor tissue and mammary gland fat pad (Cutroneo *et al*, 1973). Slow cell turnover as judged by rate of clearance of DNA-bound radioactivity (Heiniger *et al*, 1972). Harderian gland has a high

porphyrin content (Margolis, 1971). Low hepatic ammonia-lyase activity in two substrains (Hanford *et al*, 1974). Low spermatozoal beta-glucuronidase activity (Erickson, 1976). Low basal level of renal glutathione S-transferase but high basal level of renal glutathione reductase (Misra *et al*, 1991). Low hepatic nicotinamide N-methyltransferase levels (Scheller *et al*, 1996). High level of alpha-fetoprotein in plasma at seven days (Adinolfi *et al*, 1990). Resistant to the development of atherosclerosis on a semi-synthetic high fat diet (Nishina *et al*, 1993). Loci on chromosomes 1, 3, 5 and 11 are associated with variation in high density lipoprotein levels with coordinate expression of cholesterol-7-alpha hydroxylase in a cross involving atherosclerosis susceptible C57BL/6 mice (Machleder *et al*, 1997). Hepatic iodothyronine deiodinase activity was only 18% of that found in C57BL/6 mice (Schoenmakers *et al*, 1993). Decreased levels of deiodinase mRNA

and hyperthyroxinemia associated with a 21-base pair insert in the promoter region of the type 1 deiodinase gene (Maia *et al*, 1995). Resistant to severe hypercapnia with hypoxia assessed by elevated minute ventilation rate (Tankersley *et al*, 1994). Has a slow and deep breathing pattern phenotype (contrast C57BL/6) (Tankersley *et al*, 1997). High intra-ocular pressure (John *et al*, 1997).

Reproduction

Breeding performance intermediate/good. Good reproductive performance, litter size 6.4, sterility 10% (Nagasawa *et al*, 1973). Large litter size, high proportion of females produce four or more litters and high proportion of fertile matings (Fernandes *et al*, 1973). C3H/HeJ has shorter and less regular oestrus cycles than C57BL/6J (Nelson *et al*, 1992). Early opening of vagina and first cornification (compared with C57BL/6 and DBA/2), but late onset of cyclicity (Nelson *et al*, 1990).

REFERENCES

- Adinolfi M, Beck SE, Seller MJ, Fedor T, McLaren A (1990) Alpha-fetoprotein levels in different strains of mice during development. *Exp. Clin. Immunogenet* 7, 123-128.
- Adkison DL, Sundberg JP (1991) "Lipomatous" hamartomas and choristomas in inbred laboratory mice. *Vet. Pathol.* 28, 305-312.
- Akamatsu Y, Barton BP (1974) Neoplasms and amyloidosis in strains of mice treated with 3-methylcholanthrene. *J. Nat. Cancer Inst.* 52, 377.
- Albert S, Wolf PL, Pryjma I, Moore W (1965) Thymus development in high and low-leukemic mice. *J. Reticuloendothel. Soc.* 2, 218-237.
- Ammassari-Teule M, Hoffman HJ, Rossi-Arnaud C (1993) Learning in inbred mice: strain-specific abilities across three radial maze problems. *Behav. Genet.* 23, 405-412.
- Amsbaugh DF, Hansen CT, Percott B, Stashak PW, Barthold DR, Baker RJ (1972) Genetic control of the antibody response to type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness. *J. Exp. Med.* 136, 931-949.
- Bang FB, Warwick A (1960) Mouse macrophages as host cells for the mouse hepatitis virus and the genetic basis of their susceptibility. *Proc. Natl. Acad. Sci. USA* 46, 1065-1071.
- Barrett CP, Donati EJ, Volz JE, Smith EB (1975) Variations in serum calcium between strains of inbred mice. *Lab. Anim. Sci.* 25, 638-640.
- Barthold SW, Beck DS, Hansen GM, Terwilliger GA, Moody KD (1990) Lyme borreliosis in selected strains and ages of laboratory mice. *J. Infect. Dis.* 162, 133-138.
- Beamer WG, Donahue LR, Rosen CJ, Baylink DJ (1996) Genetic-variability in adult bone-density among inbred strains of mice. *Bone* 18, 397-403.
- Belyaev DK, Gruntenko EV, Vilelets IY (1970) Genetic differentiation of the thymus in mice of different strains with respect to malignant growth communication. II. Differences in the weight of the thymus in various strains of mice. *Sov. Genet.* 6, 47-51.
- Bennett LM, Farnham RJ, Drinkwater NR (1995) Strain-dependent differences in DNA synthesis and gene expression in the regenerating livers of C57BL/6J and C3H/HeJ mice. *Molecular Carcinogenesis* 14, 46-52.
- Bentvelzen P, Daams JH, Hageman P, Calafat J (1970) Genetic transmission of viruses that incite mammary tumors in mice. *Proc. Natl. Acad. Sci. USA* 67, 377-384.
- Bergman RK, Munoz J (1968) Action of the histamine sensitizing factor from *Bordetella pertussis* on inbred and random bred strains of mice. *Int. Arch. Allergy* 34, 331-338.
- Bhathal PS, Jordan TW, Mackay IR (1990) Mouse strain differences in susceptibility to sporidesmin-induced biliary tract injury. *Liver* 10, 193-204.
- Blake RL (1970) Regulation of liver tyrosine amino transferase activity in inbred strains and mutant mice. I. Strain variance in fasting enzyme levels. *Int. J. Biochem.* 1, 361-370.
- Bovet D, Bovet-Nitti F, Oliverio A (1966) Effects of nicotine on avoidance conditioning of inbred strains of mice. *Psychopharmacologia* 10, 1-5.
- Bovet D, Bovet-Nitti F, Oliverio A (1969) Genetic aspects of learning and memory in mice. *Science* 163, 139-149.
- Brown AM (1965) Pharmacogenetics of the mouse. *Lab. Anim. Care* 15, 111-118.
- Bruell JH (1964) Inheritance of behavioral and physiological characters of mice and the problem of heterosis. *Am. Zool.* 4, 125-138.
- Bruell JH (1969) Genetics and adaptive significance of emotional defecation in mice. *Ann. NY Acad. Sci.* 159, 825-830.
- Bruell JH, Daroczy AF, Hellerstein HK (1962) Strain and sex differences in serum cholesterol levels in mice. *Science* 135, 1071-1072.
- Brunnett SR (1997) Morphologic response of myocardium to freeze-thaw injury in mouse strains with dystrophic cardiac calcification. *Lab. Anim. Sci.* 47, 11-18.
- Cerny J, McAlack RF, Sajid MA, Friedman H (1971) Genetic differences in the immunocyte response of mice to separate determinants on one bacterial antigen. *Nature New Biol.* 230, 247-248.
- Chai CK, Dickie MM (1966) Endocrine variations. In: *Biology of the laboratory mouse*, 2nd ed. (Green EL, ed). New York: McGraw-Hill, pp 387-403.
- Cutroneo KR, Guzman NA, Liebelt AG (1973) Elevation of peptidylproline hydroxylase activity and collagen synthesis in spontaneous primary mammary cancers of inbred mice. *Cancer Res.* 32, 2828-2833.
- De Souza CM, Maia LCS, Vaz NM (1974) Susceptibility to cutaneous anaphylaxis in inbred strains of mice. *J. Immunol.* 112, 1369-1372.
- DeLost P, Chirvan-Nia P (1958) Differences raciales dans l'involution de la zone x multi surrenalienne chez la souris adulte vierge. *C. R. Soc. Biol.* 152, 453-455.
- Deringer MK, Dunn TB, Heston WE (1953) Results of exposure of strain C3H mice to chloroform. *Proc. Soc. Exp. Biol. Med.* 83, 474-479.
- Di Pauli R (1972) Genetics of the immune response. I. Differences in the specificity of antibodies to lipopolysaccharides among different strains of mice. *J. Immunol.* 109, 394-400.
- Dileto CL, Travis EL (1996) Fibroblast radiosensitivity in-vitro and lung fibrosis in-vivo - comparison between a fibrosis-prone and fibrosis-resistant mouse strain. *Radiation Res.* 146, 61-67.
- Dragani TA, Manenti G, Gariboldi M, De Gregorio L, Pierotti MA (1995) Genetics of liver tumor susceptibility in mice. *Toxicol. Lett.* 82-83, 613-619.
- Enomoto T, Weghorst CM, Ward JM, Anderson LM, Perantoni AO, Rice JM (1993) Low frequency of H-ras activation in naturally occurring hepatocellular tumors of C3H/HeNCR mice. *Carcinogenesis* 14, 1939-1944.
- EORTC Screening group (1972) Handbook of materials and methods. *Eur. J. Cancer* 8, 185-196.
- Erickson RP (1976) Strain variation in spermatazoal -glucuronidase in mice. *Genet. Res.* 28, 139-145.
- Evans JG, Collins MA, Lake BG, Butler WH (1992) The histology and development of hepatic nodules and carcinoma in C3H/He and C57BL/6 mice following chronic phenobarbition administration. *Toxicologic Pathology* 20, 585-594.
- Fahay JL (1965) Differences in the electrophoretic mobility of antibody from inbred strains of mice. *J. Immunol.* 94, 819-823.
- Faulkner CB, Davidson MK, Davis JK, Schoeb TR, Simecka JW, Lindsey JR (1995) Acute Mycoplasma pulmonis infection associated with coagulopathy in C3H/HeN mice. *Lab. Animal Sci.* 45, 368-372.
- Fenton PF, Dowling MT (1953) Studies on obesity. I. Nutritional obesity in mice. *J. Nutr.* 49, 319-331.
- Fernandes G, Yunis EJ, Good RA (1973) Reproductive deficiency of NZB male mice: Possibility of a viral basis. *Lab. Invest.* 29, 278-281.
- Festing MFW (1973) Water escape learning in mice. I. Strain differences and biometrical considerations. *Behav. Genet.* 3, 13-24.
- Festing MFW (1997) Inbred Strains of mice. *Mouse genome* 95, 519-686.
- Festing MFW, Blackmore DK (1971) Life span of specified-pathogen-free (MRC category 4) mice and rats. *Lab. Anim.* 5, 179-192.
- Flaks A (1968) The susceptibility of various strains of neonatal mice to the carcinogenic effects of 9, 10-dimethyl-1, 2-benzanthracene. *Eur. J. Cancer* 4, 579-585.
- Freund R, Dubensky T, Bronson R, Sotnikov A, Carroll J, Benjamin T (1992) Polyoma tumorigenesis in mice: evidence for dominant resistance and tumor susceptibility genes of the host. *Virology* 191, 724-731.
- Fuchs S, Mozes E, Maoz A, Sela M (1974) Thymus independence of a collagen-like synthetic polypeptide and of collagen, and the need for thymus and bone marrow-cell cooperation in the immune response to gelatin. *J. Exp. Med.* 139, 148-158.
- Gabridge MG, Abrams GD, Murphy WH (1972) Lethal toxicity of Mycoplasma fermentans in mice. *J. Infect. Dis.* 125, 153-160.
- Gaines S, Currie JA, Tully JG (1965) Factors affecting formation of incomplete Vi antibody in mice. *J. Bacteriol.* 90, 635-642.
- Gasser DL, Silvers WK (1971) Genetic basis of male skin rejection in mice. *Transplant* 12, 412-414.
- Green MC, Azar CA, Maren TH (1973) Strain differences in susceptibility to the teratogenic effect of acetazolamide in mice. *Teratology* 8, 143-145.
- Hanford WC, Nep RL, Arlin SM (1974) Genetic variation in histidine ammonia-lyase activity in the mouse. *Biochem. Biophys. Res. Comm.* 61, 1434-1437.
- Harada M, Nagata M, Takeuchi M, Ohara T, Makino S, Watanabe A (1991) Age-dependent difference in susceptibility to IgE antibody- and IgG1 antibody-mediated passive anaphylactic shock in the mouse. *Immunological Investigations* 20, 515-523.
- Haston CK, Amos CI, King TM, Travis EL (1996) Inheritance of susceptibility to bleomycin-induced pulmonary fibrosis in the mouse. *Cancer Res.* 56, 2596-2601.
- Hegmann JP (1972) Physiological function and behavioral genetics. I. Genetic variance for peripheral nerve conduction velocity in mice. *Behav. Genet.* 2, 55-67.
- Heiniger HJ, Chen HW, Meier H, Taylor BA, Commerford LS (1972) Studies on the genetic control of cell proliferation. I. Clearance of DNA-bound radioactivity in 19 inbred strains and hybrid mice. *Life Sci.* 11, 87-98.
- Heiniger HJ, Taylor BA, Hards EJ, Meier H (1975) Heritability of the phytohemagglutinin responsiveness of lymphocytes and its relationship to leukemogenesis. *Cancer Res.* 35, 825-831.
- Hellman A, Fowler AK (1972) Studies of the blastogenic response of murine lymphocyte. III. Specific viral transformation. *Proc. Soc. Exp. Biol. Med.* 141, 106-109.
- Henderson ND (1970) Genetic influences on the behavior of mice can be obscured by laboratory rearing. *J. Comp. Physiol. Psychol.* 72, 505-511.
- Heston WE (1963) Genetics of neoplasia. In: *Methodology in mammalian genetics* (Burdette WJ, ed). San Francisco: Holden-Day, pp 247-268.
- Heston WE, Vlahakis G (1971) Mammary tumors, plaques and hyperplastic alveolar nodules in various combinations of mouse inbred strains and the different lines of the mammary tumor virus. *Int. J. Cancer* 7, 141-148.
- Hill GB, Osterhout S, O'Fallon WM (1968) Variation in response to hyperbaric oxygen among inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 129, 687-689.
- Hoag WG (1963) Spontaneous cancer in mice. *Ann. NY Acad. Sci.* 108, 805-831.
- Hoffman HA, Rechcigl M Jr (1971) Erythrocyte catalase in inbred mice. *Enzyme* 12, 219-225.
- Hummel KP, Richardson FL, Fekete E (1966) Anatomy. In *Biology of the Laboratory Mouse*, 2nd ed. (Green EL, ed). New York: McGraw-Hill, pp 247-307.
- Ingram DK, Reynolds MA, Les EP (1982) The relationship of genotype, sex, body weight, and growth parameters to lifespan in inbred and hybrid mice. *Mech. Ageing Dev.* 20, 253-266.
- James K, Milne I (1972) The effect of anti-lymphocytic antibody on the humoral immune response in different strains of mice. I. The response to bovine serum albumin. *Immunol.* 23, 897-909.
- Jenson AB, Groff DE, McConahy RJ, Dixon FJ (1976) Transmission of murine leukemia virus (Scripps) from parent to progeny mice as determined by P30 antigenemia. *Cancer Res.* 36, 1228-1232.
- John SWM, Hagaman JR, MacTaggart TE, Peng L, Smithes O (1997) Intraocular pressure in inbred mouse strains. *Investigative Ophthalmology & Visual Science* 38, 249-253.
- Johnson AP, Tuffrey M, Taylor-Robinson D (1989) Resistance of mice to genital infection with *Neisseria gonorrhoea*. *J. Med. Microbiol.* 30, 33-36.
- Kallis N (1972) Transplanted tumors. *Jax Notes* 410, Jackson Laboratory, Bar Harbor, Maine.
- Kano K, Mizuma Y (1974) Comparison of the total blood volume in four inbred strains of mice with different hemoglobin types. *Exp. Animals (Japan)* 23, 123-127.
- Kawashima I, Nakamura O, Tai T (1992) Antibody responses to ganglioside series gangliosides in different strains of inbred mice. *Molecular Immunology* 29, 625-632.
- Kleeberger SR, Levitt RC, Zhang LY (1993) Susceptibility to ozone-induced inflammation. I. Genetic control of the response to subacute exposure. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 264, L15-L20.
- Konno S, Adachi M, Matsuura T, Sunouchi K, Hoshino H, Okazawa A, Kobayashi H, Takahashi T (1993) Bronchial reactivity to methacholine and serotonin in six inbred mouse strains. [Japanese]. *Japanese Journal of Allergy* 42, 42-47.
- Kouri RE, Salerno RA, Whitmore CE (1973) Relationships between arylhydrocarbon hydroxylase inducibility and sensitivity to chemically induced subcutaneous sarcomas in various strains of mice. *J. Natl. Cancer Inst.* 50, 363-368.
- Law LW (1966) Studies of thymic function with emphasis on the role of the thymus in oncogenesis. *Cancer Res.* 26, 551-574.
- Les EP (1972) A disease related to cage population density: tail lesions of C3H/HeJ mice. *Lab. Animal Sci.* 22, 56-60.
- Levine BB, Vaz NM (1970) Effect of combinations of inbred strain, antigen and antigen dose on immune responsiveness and reagin production in the mouse. *Int. Arch. Allergy* 39, 156-171.
- Liebelt AG, Liebelt RA, Dmochowski L (1971) Cytoplasmic inclusion bodies in primary and transplanted hepatomas of mice of different strains. *J. Natl. Cancer Inst.* 47, 413-427.
- Liebelt RA, Suzuki S, Liebelt AG, Lane M (1970) Virus-like particles in chemically induced sarcomas in high- and low-leukemia strains of mice. *Cancer Res.* 30, 2438-2448.
- Lieberman J, Kellog F (1967) Hyaline-membrane formation and pulmonary plasminogen-activator activity in various strains of mice. *Pediatrics* 39, 75-81.
- Lush IE, Arnold CJ (1975) High coumarin 7-hydroxylase activity does not protect mice against Warfarin. *Heredity* 35, 279-281.
- Lyon MF, Rastan S, Brown SDM (1996) Genetic variants and strains of the laboratory mouse. 2 Volumes. Oxford, New York, Tokyo: Oxford University Press.
- Machleder D, Ivandic B, Welch C, Castellani L, Reue K, Lusk AJ (1997) Complex genetic control of HDL levels in mice in response to an atherogenic diet - Coordinate regulation of HDL levels and bile acid metabolism. *J. Clin. Invest.* 99, 1406-1419.
- Maia AL, Berry MJ, Sabbag R, Hamey JW, Larsen PR (1995) Structural and functional differences in the diol gene in mice with inherited type 1 diolase deficiency. *Molec. Endocrinol.* 9, 969-980.
- Margolis FL (1971) Regulation of porphyrin biosynthesis in the Harderian gland of inbred mouse strains. *Arch. Biochem. Biophys.* 145, 373-381.
- Marks MJ, Stitzel JA, Collins AC (1989) Genetic influences on nicotine responses. *Pharmacol. Biochem. Behav.* 33, 667-678.
- Matylniak JE, Reiner SL (1995) T helper phenotype and genetic susceptibility in experimental Lyme disease. *Journal Of Experimental Medicine* 181, 1251-1254.
- McCarthy MM, Dutton RW (1975) The humoral response of mouse spleen cells to two types of sheep erythrocytes. *J. Immunol.* 115, 1316-1321.

90. McClearn GE, Wilson JR, Meredith W (1970) The use of isogenic and heterogenic mouse stocks in behavioral research. In: Contribution to behavior genetic analysis. The mouse as a prototype (Lindzey G, Thiessen DD, eds). New York: Appleton-Century-Crofts, pp 3-32.
91. McLaren A, Tait A (1969) Cytoplasmic isocitrate dehydrogenase variation within the C3H inbred strain. *Genet. Res.* 14, 93.
92. Misra M, Rodriguez RE, North SL, Kasprzak KS (1991) Nickel-induced renal lipid peroxidation in different strains of mice: concurrence with nickel effect on antioxidant defence systems [published erratum appears in *Toxicol. Lett.* (1992) 60, 239]. *Toxicol. Lett.* 58, 121-133.
93. Muhlbock O, Tengbergen WP Jr (1971) Instability of characteristics in inbred strains of mice. In: Defining the laboratory animal (Schneider HA, ed), Proc. IV ICLAS Symposium, pp 230-249.
94. Murray WS, Little CC (1967) Genetic studies of carcinogenesis in mice. *J. Natl. Cancer Inst.* 38, 639-656.
95. Nagasawa H, Miyamoto M, Fujimoto M (1973) Reproductivity in inbred strains of mice and project for their efficient production. *Exp. Animals (Japan)* 22, 119-126.
96. Nagy ZM, Misanin JR (1970) Visual perception in the retinal degenerate C3H mouse. *J. Comp. Physiol. Psychol.* 72, 306-310.
97. Nandi J, Bern HA, Biglieri EG, Pieprzak JK (1967) In vitro steroidogenesis by the adrenal glands of mice. *Endocrinol.* 80, 576-582.
98. Neal RA, Harris WG (1975) Attempts to infect inbred strains of rats and mice with *Entamoeba histolytica*. *Trans. R. Soc. Trop. Med. Hyg.* 69, 429-430.
99. Nelson JF, Karelus K, Felicio LS, Johnson TE (1990) Genetic influences on the timing of puberty in mice. *Biol. Reprod.* 42, 649-655.
100. Nelson JF, Karelus K, Felicio LS, Johnson TE (1992) Genetic influences on oestrous cyclicity in mice: evidence that cycle length and frequency are differentially regulated. *J. Reprod. Fertil.* 94, 261-268.
101. Niewiesk S, Brinckmann U, Bankamp B, Sirak S, Liebert UG, Ter Meulen V (1993) Susceptibility to measles virus-induced encephalitis in mice correlates with impaired antigen presentation to cytotoxic T lymphocytes. *Journal of Virology* 67, 75-81.
102. Nikulina EM, Skirinskaya JA, Popova NK (1991) Role of genotype and dopamine receptors in behavior of inbred mice in a forced swimming test. *Psychopharmacology* 105, 525-529.
103. Nishina PM, Wang J, Toyofuku W, Kuypers FA, Ishida BY, Paigen B (1993) Atherosclerosis and plasma and liver lipids in nine inbred strains of mice. *Lipids* 28, 599-605.
104. Nishioka Y (1987) Y-chromosomal DNA polymorphism in mouse inbred strains. *Genet. Res.* 50, 69-72.
105. Oesch F, Morris N, Daly JW (1973) Genetic expression of the induction of epoxide hydrase and aryl hydrocarbon hydroxylase activities in the mouse by phenobarbital or 3-methylcholanthrene. *Molec. Pharmacol.* 9, 692-696.
106. Paul WE, Yoshida T, Benacerraf B (1970) Genetic control of the specificity of anti-DNP antibodies. II. Differences in the specificity of anti-DNP antibody produced by several inbred strains of mice. *J. Immunol.* 105, 314-321.
107. Peacock AC, Gelderman AH, Ragland RH, Hoffman HA (1967) Haptoglobin levels in serum of various strains of mice. *Science* 158, 1703-1704.
108. Pennycook PR (1967) The comparison of the effects of a variety of factors on the metabolic rate of the mouse. *Aus. J. Bio. Med. Sci.* 45, 331-346.
109. Petty RE, Steward MW, Soothill JF (1972) The heterogeneity of antibody affinity in inbred mice and its possible immunopathologic significance. *Clin. Exp. Immunol.* 12, 231-241.
110. Pinchuck P, Maurer PH (1965) Antigenicity of polypeptides (poly alpha amino acids). XVI. Genetic control of immunogenicity of synthetic polypeptides in mice. *J. Exp. Med.* 122, 673-679.
111. Plant J, Glynn AA (1974) Natural resistance to Salmonella infection, delayed hypersensitivity and Ir genes in different strains of mice. *Nature* 248, 345-347.
112. Pope BL, Chourmouzis E, MacIntyre JP, Lee S, Goodman MG (1994) Murine strain variation in the natural killer cell and proliferative responses to the immunostimulatory compound 7-Allyl-8-oxoguanosine: Role of cytokines. *Cell. Immunol.* 159, 194-210.
113. Price P, Eddy KS, Papadimitriou JM, Faulkner DL, Shellam GR (1991) Genetic determination of cytomegalovirus-induced and age-related cardiopathy in inbred mice. Characterization of infiltrating cells. *Am. J. Pathol.* 138, 59-67.
114. Qiao JH, Fishbein MC, Demer LL, Lusic AJ (1995) Genetic determination of cartilaginous metaplasia in mouse aorta. Atherosclerosis, Thrombosis and Vascular Biology 15, 2265-2272.
115. Rager-Zisman B, Ju G, Udem S (1976) Resistance and susceptibility of mice to infection with measles virus. *Fed. Proc.* 35, 391.
116. Ram JS, Dehellis RA, Glenner GG (1969) Amyloid. VIII on strain variation in experimental murine amyloidosis. *Proc. Soc. Exp. Biol. Med.* 130, 462-464. Rank WR, Flugge U, DiPauli R (1969) Inheritance of the lipid A- induced 19S-plaque-forming cell-response in mice. Evidence for three antigen-recognition mechanisms. *Behringwerk-Mitt.* 49, 222-229.
117. Richardson FL, Hall G (1963) Mammary tumors, mammary gland development and morphology of ovaries in virgin mice of strains C3H, CBA and their F1 hybrids. *J. Natl. Cancer Res.* 31, 529-539.
118. Riley V (1975) Mouse mammary tumors: alteration of incidence as an apparent function of stress. *Science* 189, 465-467.
119. Robinson SF, Marks MJ, Collins AC (1996) Inbred mouse strains vary in oral self-selection of nicotine. *Psychopharmacology* 124, 332-339.
120. Roderick TH (1963) The response of twenty-seven inbred strains of mice to daily doses of whole-body X-irradiation. *Radiation Res.* 20, 631-639.
121. Rubinstein P, Liu N, Strenn EW, Decary F (1974) Electrophoretic mobility and agglutinability of red blood cells: a new polymorphism in mice. *J. Exp. Med.* 139, 313-322.
122. Russell ES, Neufeld EF, Higgins CT (1951) Comparison of normal blood picture of young adults from 18 inbred strains of mice. *Proc. Soc. Exp. Biol. Med.* 78, 761-766.
123. Sakagami T, Dixon M, O'Rourke J, Howlett R, Alderuccio F, Vella J, Shimoyama T, Lee A (1996) Atrophic gastric changes in both *Helicobacter felis* and *Helicobacter pylori* infected mice are host dependent and separate from antral gastritis. *Gut* 39, 639-648.
124. Sato H, Nagayoshi M, Ikawa Y, Kato MV (1996) A C3H strain-specific allele (alpha val26) of the murine alpha-globin gene newly detected by UT-PAGE and RT-PCR-SSCP analysis. *Mutat. Res.* 351, 125-132.
125. Scheller T, Orgacka H, Szumlanski CL, Weinshilboum RM (1996) Mouse liver nicotinamide N-methyltransferase pharmacogenetics: Biochemical properties and variation in activity among inbred strains. *Pharmacogenetics* 6, 43-53.
126. Schlager G, Dickie MM (1967) Spontaneous mutations and mutation rates in the house mouse. *Genetics* 57, 319-330.
127. Schoenmakers CHH, Pigmanis IGAJ, Poland A, Visser TJ (1993) Impairment of the selenoenzyme type I iodothyronine deiodinase in C3H/He mice. *Endocrinol.* 132, 357-361.
128. Scott P, Eaton A, Gause WC, Zhou XD, Hondowicz B (1996) Early IL-4 production does not predict susceptibility to *Leishmania major*. *Experimental Parasitology* 84, 178-187.
129. Shaikh ZA, Jordan SA, Tewari PC (1993) Cadmium disposition and metallothionein induction in mice: Strain-, sex-, age- and dose-dependent differences. *Toxicology* 80, 51-70.
130. Shepard CC, Habas JA (1967) Relation of infection to tissue temperature in mice infected with *Mycobacterium marinum* and *Mycobacterium leprae*. *J. Bacteriol.* 93, 790-796.
131. Shih DM, Gu L, Hama S, Xia YR, Navab M, Fogelman AM, Lusic AJ (1996) Genetic-dietary regulation of serum paraoxonase expression and its role in atherosclerosis in a mouse model. *J. Clin. Invest.* 97, 1630-1639.
132. Shimosato K, Saito T, Marley RJ (1994) Genotype-specific blockade of cocaine-induced weight loss by the protein synthesis inhibitor, anisomycin. *Life Sciences* 55, PL293-PL299.
133. Sinha YM, Salocks CB, Vanderlaan WP (1975) Prolactin and growth hormone levels in different inbred strains of mice: patterns in association with estrous cycle, time of day and perphenazine stimulation. *Endocrinol.* 97, 1112-1122.
134. Smith AM (1976) The effects of age on the immune response to type III pneumococcal polysaccharide (SIII) and bacterial lipopolysaccharide (LPS) in BALB/c, SJL/J and C3H mice. *J. Immunol.* 116, 469-474.
135. Smith GS, Walford RL (1977) Influence of the main histocompatibility complex on ageing in mice. *Nature* 270, 727-729.
136. Smith GS, Walford RL (1978) Influence of the H-2 and H-1 histocompatibility systems upon life-span and spontaneous cancer incidence in congenic mice. *Birth defects: Original article series* 14, 281-312.
137. Southwick CH, Clark LH (1966) Aggressive behavior and exploratory activity in fourteen mouse strains. *Am. Zool.* 6, 559.
138. Southwick CH, Clark LH (1968) Interstrain differences in aggressive behavior and exploratory activity of inbred mice. *Commun. Behav. Biol. Part A* 1, 49-59.
139. Srivastava RAK (1995) Increased apoB100 mRNA in inbred strains of mice by estrogen is caused by decreased RNA editing protein mRNA. *Biochemical and Biophysical Research Communications* 212, 381-387.
140. Stasik JH (1970) Inheritance of T-maze learning in mice. *J. Comp. Physiol. Psychol.* 71, 251-257. Stravrovadi P, Toliou T, Polyzonis MB (1991) A new approach combining interferon and bromocriptine LA to prevent spontaneous mouse mammary carcinoma. *Journal of Interferon Research* 11, 49-51.
141. Strong LC (1952) Differences in response among mice of fifteen inbred strains to the subcutaneous injection of methylcholanthrene. *Yale J. Biol. Med.* 25, 34-43.
142. Takahashi M, Kleeberger SR, Croxton TL (1995) Genetic control of susceptibility to ozone-induced changes in mouse tracheal electrophysiology. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 269, L6-110.
143. Tanioka Y, Esaki K (1971) Strain differences in mortality of anaphylactic shock in mice-challenging by intravenous injection. *Exp. Animals (Japan)* 20, 127-130.
144. Tankersley CG, Fitzgerald RD, Kleeberger SR (1994) Differential control of ventilation among inbred strains of mice. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology* 267, R1371-R1377.
145. Tankersley CG, Fitzgerald RD, Levitt RC, Mitzner WA, Ewart SL, Kleeberger SR (1997) Genetic control of differential baseline breathing pattern. *J. Appl. Physiol.* 82, 874-881.
146. Taylor BA (1976) Genetic analysis of susceptibility to isoniazid-induced seizures in mice. *Genetics* 83, 373-377.
147. Tengbergen WJPR (1970) Morphological classification of mammary tumors in the mouse. *Path. Eur.* 5, 260-272.
148. Thomas PE, Hutton JJ, Taylor BA (1973) Genetic relationship between aryl hydrocarbon hydroxylase inducibility and chemical carcinogen induced skin ulceration in mice. *Genetics* 74, 655-659.
149. Thompson WR (1953) The inheritance of behavior: behavioral differences in fifteen mouse strains. *Can. J. Psychol.* 7, 145-155.
150. Toda S, Kimura M, Tohya K (1989) Strain differences in histamine release from mouse peritoneal mast cells induced by compound 48/80 or A23187. *Jikken Dobutsu - Experimental Animals* 38, 135-137.
151. Tokunaga Y, Hiramine C, Itoh M, Mukasa A, Hojo K (1993) Genetic susceptibility to the induction of experimental allergic encephalomyelitis (EAO) without adjuvant. I. Comparison of pathology, delayed type hypersensitivity, and antibody. *Clin. Immunol. Immunopathol.* 66, 239-247.
152. Tsubura A, Senzaki H, Oyaizu T, Fujita Y, Morii S (1993) Strain differences in neoplastic response to DMBA-induced uterine vascular tumors in mice. *International Journal of Oncology* 2, 927-930.
153. Valatx JL, Bugat R (1974) Facteurs génétiques dans le déterminisme du cycle veille-sommeil chez la souris. *Brain Res.* 69, 315-330.
154. Vaz NM, Phillips-Quagliata JM, Levine BB, Vaz EM (1971) H-2 linked genetic control of immune responsiveness to ovalbumin and ovomucoid. *J. Exp. Med.* 134, 1335-1348.
155. Vesell ES (1968) Factors altering the responsiveness of mice to hexobarbital. *Pharmacology* 1, 81-97.
156. Wetzel BK, Wolman SH (1969) Fine structure of a second kind of thyroid follicle in the C3H mouse. *Endocrinology* 84, 563-578.
157. Whitmire CE, Salerno RA (1972) RNA tumour virus antigen and tumour induction by various doses of 3-methylcholanthrene in various strains of mice treated as weanlings. *Cancer Res.* 32, 1129-1132.
158. Whitmire CE, Salerno RA, Rabstein LS, Heubner RJ, Turner HC (1971) RNA tumor-virus antigen expression in chemically induced tumors. Virus-genome specified common antigens detected by complement fixation in mouse tumors induced by 3-methylcholanthrene. *J. Natl. Cancer Inst.* 47, 1255-1265.
159. Whitmore AC, Whitmore SP (1985) Subline divergence within Strong's C3H and CBA inbred strains. A review. *Immunogenetics* 21, 407-428.
160. Williams NM, Timoney PJ (1994) Variation in susceptibility of ten mouse strains to infection with a strain of *Ehrlichia risticii*. *J. Comp. Pathol.* 110, 137-143.
161. Yamamoto Y, Saito H, Setogawa T, Tomioka H (1991) Sex differences in host resistance to *Mycobacterium marinum* infection in mice. *Infect. Immun.* 59, 4089-4096.
162. Young CR, Deacon NJ, Ebringer A, Davis DAL (1976) Genetic control of the immune response to ferritin in mice. *J. Immunogenet.* 3, 199-205.
163. Zhang LY, Levitt RC, Kleeberger SR (1995) Differential susceptibility to ozone-induced airways hyperreactivity in inbred strains of mice. *Experimental Lung Research* 21, 503-518.