

SHR (Spontaneous Hypertensive Rat)

SHR/NHsd

Okamoto and Aoki, 1963, from an outbred Wistar Kyoto male with spontaneous hypertension and a female with slightly higher than normal blood pressure. Subsequently brother x sister pairs were selected for spontaneous hypertension (Okamoto, 1969; Okamoto *et al*, 1972). A number of sublines have been developed with a tendency to develop cardiovascular lesions and stroke (see particularly SHRSP) (Nagaoka *et al*, 1976), and hypercholesterolemia (Yamori, 1984). For a more recent review; see Yamori, (1994). From Department of Pathology, Faculty of Medicine, Kyoto University to National Institutes of Health, Bethesda in 1966.

From National Institutes of Health to Harlan Laboratories. Harlan became Envigo in 2015, then Envigo was acquired by Inotiv in 2021.

CHARACTERISTICS

Animal model

SHR rats express insulin resistance, and are a suitable model for insulin resistance and essential hypertension in non-obese humans (Swislocki and Tsuzuki, 1993). SHR rats are hyperactive and may be a useful model for childhood hyperkinesia and attention-deficit hyperactivity disorder (Sagvolden *et al*, 1993). SHR rats have been used as a model for screening anti-hypertensive drugs (Roba, 1976).

Anatomy

Low ten week body weight in males (Tanase *et al*, 1982). High relative heart weight in ten week old males (Tanase *et al*, 1982). SHR rats express insulin resistance, and are a suitable model for insulin resistance and essential hypertension in non-obese humans (Swislocki and Tsuzuki, 1993). SHR rats have fewer glomeruli than WKY rats, but they are of similar size, resulting in a reduced glomerular volume. This is consistent with the hypothesis that the kidney plays an important role in hypertension (Skov *et al*, 1994). Fetal but not placental weight is reduced compared with WKY (Johnston, 1995).

Behavior

SHR rats are hyperactive and may be a useful model for childhood hyperkinesia and attention-deficit hyperactivity disorder (Sagvolden *et al*, 1993).

Open field exploratory behavior was greater in SHR than WKY rats, and the most active SHR developed the most pronounced hypertension (Knardahl and Sagvolden, 1979). Muricidal behavior of 90% was reported (Roba, 1976).

Drugs

Genetically resistant to the induction of mammary tumors by dimethylbenz(a)anthracene due to a blockade of tumor promotion (Harris *et al*, 1994). Roba (1976) concluded that the strain is a suitable model for screening anti-hypertensive drugs.

Genetics

Cross analysis between SHR and Donryu rats indicated that the hypertensive trait of SHR is regulated by a single gene and several other genes with minor effect (Tanase, 1979; Roba, 1976). Esterase alleles have been described by Womack (1973). There is no evidence for substrain differentiation among SHR stocks from the major commercial suppliers in the USA both respect to phenotype and DNA fingerprints (Blizard *et al*, 1991).

Coat color genes

- *c* : albino.

Histocompatibility

- *RT1^k*

Biochemical markers

- *Es-1^a, Es-2^a, Es-3^b, Es-4^a, Gc^a, Hbb^a, Pep-3^a, Pgd^b*

Husbandry

The "Committee on the care and use of hypertensive rats (1976)" has issued guidelines for the breeding and care of this strain.

Life-span and spontaneous disease

Histopathologic changes appeared in males when they became eight months old, in females at an age of 12-15 months. Female SHR rats live significantly longer than males (Wexler *et al*, 1981).

A few cases of multiple osteosarcoma in the skull, caudal vertebrae, and fore- and hindlimbs were reported (Iida *et al*, 1979). Periarteritis nodosa, which starts at level of duodenal artery and proceeds posteriorly to the testicular arteries, is the most common lesion. Myocardial lesions and nephrosclerosis are common; cerebral lesions are usually seen in rats with blood pressure greater than 200 mm Hg (Roba, 1976).





High blood pressure, reaching 171 mmHg at ten weeks of age (Tanase *et al*, 1982). Blood pressure of about 200 mm in mature males and 175 mm in mature females (Roba, 1976). According to Yamori (1984), the rats develop hypertension spontaneously without exception at the age of 7-15 weeks. There is a systolic blood pressure plateau of about 200 mmHg. The genetic basis is polygenic, with at least three major genes involved (Tanase and Suzuki, 1971; Yen *et al*, 1974).

There is a high incidence of cardiovascular disease (Okamoto *et al* 1973), but a low incidence of stroke, which can be increased to about 30% with chronic stress (Yamori, 1984). Alloxan diabetes further increases blood pressure, but the animals respond to anti-hypertensive drugs (Okamoto, 1969). Yamori states that SHR rats show a functional increase in peripheral vascular resistance, which mostly depend on neurogenic mechanisms, which probably originate in a disorder of central blood pressure regulation. The blood pressure per se and increased neurogenic tone accelerate cardiovascular protein synthesis and induce structural vascular changes, which contribute to the maintenance of the hypertension. Studies on cultured vascular smooth muscle suggest a genetic predisposition to hyperplastic growth of these cells and its stimulation by β -adrenergic mechanisms. According to Dietz *et al* (1984) there is an abnormality of intracellular electrolyte balance with increased intracellular sodium and calcium concentration. Grobecker *et al* (1975) found that in young SHR rats the plasma levels of both noradrenaline and dopamine- β -hydroxylase were increased over control WKY rats, but total catecholamines were not significantly different. Catecholamine

content of the adrenals was reduced. Circulating thyrotrophin levels were markedly elevated over two control strains (Werner *et al*, 1975). There was a reduced ^{131}I metabolism and increased thyroid weight relative to Wistar controls (Fregley, 1975).

Environmental and dietary factors can influence the degree of hypertension (Yamori *et al* 1978; Yamori *et al* 1986). A high (8%) salt diet increased systolic blood pressure, but not so much as in strain SS/Jr (Adams and Blizard, 1991). A polymorphism in the heat shock protein 70 (hsp70) mapping in the RT1 complex was found to be associated with variation in blood pressure of 15 mm Hg among recombinant inbred strains (Hamet *et al*, 1992). Strain is significantly more sensitive to the hypotensive effects of GABA than normotensive Sprague Dawley or WKY rats, with evidence that the effects are mediated by the brain angiotensin system (Roberts *et al*, 1993). Plasma renin and angiotensin II levels are not elevated (Cambell *et al*, 1995). Glucose turnover in lean and obese (carrying the fatty gene) SHR rats has been described by Berdanier *et al*, (1993). There is reduced cancellous bone mass in SHR compared with WKY (Wang *et al*, 1993). The Y-chromosome of SHR increases blood pressure when backcrossed to strain WKY for 11 generations (Ely *et al*, 1993). There is a deficit in visual acuity at 40-66 days, prior to the onset of hypertension, and it is particularly marked in the blue spectrum (Rogers *et al*, 1993).

Miscellaneous

Characteristics of the SHR strain have been described by Festing (1979), De Jong (1984) and Greenhouse *et al* (1990).

Nutrition

High dietary calcium (2.5%) attenuated the time course of hypertension (Ayachi, 1979). Reduction of dietary vitamin E prevents the development of hypertension, possibly due to a significant increase in prostaglandin catabolism (Pace-Asciak and Carrara, 1979). Sodium chloride solutions were preferred to deionized water (Roba, 1976; Forman and Falk, 1979). Caloric restriction without sodium restriction reduces blood pressure (Young *et al*, 1978).

Physiology and biochemistry

Renal vascular resistance is elevated (Fink and Brody, 1979; Arendshorst and Beierwaltes, 1979a). Renal function studies indicate that kidneys of SHR require a higher arterial pressure than kidneys of WKY to excrete a given amount of salt and water (Arendshorst and Beierwaltes, 1979b; Beierwaltes and Arendshorst, 1978). Renal blood flow autoregulation was as efficient as in WKY rats (Arendshorst, 1979). PGE2 synthesis in aortas and PGE2Q in inferior cava walls were increased (Limas and Limas, 1977). PGE2 synthesis in renomedullary microsomes was increased (Limas and Limas, 1979).

Reproduction

Long gestation period: $22.55 \pm .31$ days (Peters, 1986).



REFERENCES

- Adams N and Blizard DA (1991) Genetic and maternal influences in rat models of spontaneous and salt-induced hypertension. *Development Psychobiology*, 24, 507-519.
- Arendshorst WJ (1979) Autoregulation of renal blood flow in spontaneously hypertensive rats. *Circ. Res.* 44, 344-349.
- Arendshorst WJ and Beierwaltes WH (1979a) Renal and nephron hemodynamics in spontaneously hypertensive rats. *Am. J. Physiol.* 236, F246-251.
- Arendshorst WJ and Beierwaltes WH (1979b) Renal tubular reabsorption in spontaneously hypertensive rats. *Am. J. Physiol.* 237, F38-47.
- Ayachi S (1979) Increased dietary calcium lowers blood pressure in the spontaneously hypertensive rat. *Metabolism* 28, 1234-1238.
- Beierwaltes WH and Arendshorst WJ (1978) Renal function of conscious spontaneously hypertensive rats. *Circ. Res.* 42, 721-726.
- Berdanier CD, Pan JS, Hartle DK and Michaelis OE (1993) Glucose-turnover in lean and obese rats of the SHE/N-cp and LA/N-cp strains. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology* 106, 87-94.
- Blizard DA, Peterson WN and Adams N (1991) Dietary salt and accelerated hypertension - lack of subline differentiation in Spontaneously Hypertensive Rat stocks from the United-States. *J. Hypertens.* 9, 1169-1175.
- Campbell DJ, Duncan AM, Kladis A and Harrap SB (1995) Angiotensin peptides in spontaneously hypertensive and normotensive donryu rats. *Hypertension* 25, 928-934.
- Committee on the Care and Use of Spontaneously Hypertensive (SHR) Rats (1976) Spontaneously hypertensive (SHR) rats: guidelines for breeding care and use. *ILAR News* 19, G1-G20.
- De Jong W, ed (1984) *Handbook of Hypertension, Vol. 4 Experimental and genetic models of hypertension*. Amsterdam, New York, Oxford: Elsevier.
- Dietz R, Shomig A and Rascher W (1984) Pathophysiological aspects of genetically-determined hypertension in rats with special emphasis on stroke prone spontaneously hypertensive rats. In: *Handbook of Hypertension Vol. 4. Experimental and genetic models of hypertension* (de Jong W, ed). Amsterdam, New York, Oxford: Elsevier, pp. 256-285.
- Ely DL, Daneshvar H, Turner ME, Johnson ML and Salisbury RL (1993) The hypertensive Y-chromosome elevates blood-pressure in F11 normotensive rats. *Hypertension* 21, 1071-1075.
- Festing MFW (1979) Inbred strains. In: *The Laboratory Rat. Vol. I. Biology and Diseases* (Baker HJ, Lindsey JR and Weisbroth SH, eds). New York: Academic Press, pp 55-72.
- Fink GD and Brody MJ (1979) Renal vascular resistance and reactivity in the spontaneously hypertensive rat. *Am. J. Physiol.* 237, F128-F132.
- Forman S and Falk JL (1979) NaCl solution ingestion in genetic (SHR) and aortic-ligation hypertension. *Physiol. Behav.* 22, 371-377.
- Fregly MJ (1975) Thyroid activity of spontaneous hypertensive rats. *Proc. Soc. Exp. Biol. Med.* 149, 124-132.
- Greenhouse DD, Festing MFW, Hasan S and Cohen AL (1990) Catalogue of inbred strains of rats. In: *Genetic monitoring of inbred strains of rats* (Hedrich HJ, ed). Stuttgart, New York: Gustav Fischer Verlag, pp 410-480.
- Grobecker H, Roisen MF, Weise V, Saavedra JM and Kopin IJ (1975) Sympathoadrenal medullary activity in young, spontaneously hypertensive rats. *Nature* 258, 267.
- Hamet P, Kong D, Pravenec M, Kunes J, Kren V, Kir P, Sun YL and Tremblay J (1992) Restriction-fragment-length-polymorphism of hsp70-gene, localized in the RT1-complex, is associated with hypertension in spontaneously hypertensive rats. *Hypertension* 19, 611-614.
- Harris SR, Mehta RS, Hartle DK, Broderon JR and Bunce OR (1994) Failure of high-fat diets to promote mammary cancers in Spontaneously Hypertensive Rats. *Cancer Lett.* 87, 9-15.
- Iida M, Utsumi K, Ishikawa T, et al (1979) Multiple osteosarcoma in spontaneously hypertensive rats. *Exp. Anim.* 28, 497-506.
- Iwase M, Yamamoto M, Iino K, Ichikawa K, Shinohara N, Yoshinari M and Fujishima M (1998) Obesity induced by neonatal monosodium glutamate treatment in spontaneously hypertensive rats; an animal model of multiple risk factors. *Hypertens. Res.* 21, 1-6.
- Johnston BM (1995) Fetal growth-retardation and increased placental weight in the Spontaneously Hypertensive Rat. *Reproduction Fertility and Development* 7, 639-645.
- Knardahl S and Sagvolden T (1979) Open-field behavior of spontaneously hypertensive rats. *Behav. Biol.* 27, 187-200.
- Limas CJ and Limas C (1977) Vascular prostaglandin synthesis in the spontaneously hypertensive rat. *Am. J. Physiol.* 233, H493-499.
- Limas CJ and Limas C (1979) Enhanced renomedullary prostaglandin synthesis in spontaneously hypertensive rats: Role of a phospholipase A2. *Am. J. Physiol.* 236, H65-72.
- Nagaoka A, Iwatsuka H, Suzuoki A and Oksmoto K (1976) Genetic predisposition to stroke in spontaneously hypertensive rats. *Am. J. Physiol.* 230, 1354-1359.
- Okamoto K (1969) Spontaneous hypertension in rats. *Int. Rev. Exp. Pathol.* 7, 227-270.
- Okamoto K, Yamori Y, Goshima A, Park C, Haebara H, Matsumoto M, Tanaka T, Okuda T, Hazama F and Kyogoku M (1972) Establishment of the inbred strain of the spontaneously hypertensive rat and genetic factors involved in hypertension. In: *Spontaneous hypertension-its pathogenesis and complications*. (Okamoto K, ed). Tokyo: Igaku Shoin Ltd, pp. 1-8.
- Okamoto K, Yamori Y, Nosaka S, Ooshima A and Hazama F (1973) Studies on hypertension in spontaneously hypertensive rats. *Clin. Sci. Molec. Med.* 45, 11s-14s.
- Pace-Asciak CR and Carrara MC (1979) Reduction in dietary vitamin E prevents onset of hypertension in developing spontaneously hypertensive rats. *Experientia* 35, 1561-1562.
- Peters AG (1986) Length of gestation period on eight inbred strains and three outbred strains of rats. *Animal Technology* 37, 109-112.
- Roba JL (1976) The use of spontaneously hypertensive rats for the study of anti-depressive agents. *Lab. Anim. Sci.* 26, 305-319.
- Roberts KA, Wright JW and Harding JW (1993) GABA and bicuculline-induced blood-pressure changes in Spontaneously Hypertensive Rats. *J. Cardiovasc. Pharmacol.* 21, 156-162.
- Rogers LJ, Bolden S, Patrech AS and Ehrlich D (1993) Visual dysfunction in the Spontaneously Hypertensive Rat. *Physiol. Behav.* 54, 903-907.
- Sagvolden T, Petterson MB and Larsen MC (1993) Spontaneously Hypertensive Rats (SHR) as a putative animal-model of childhood hyperkinesia - SHR behavior compared to 4 other rat strains. *Physiol. Behav.* 54, 1047-1055.
- Skov K, Nyengaard JR, Korsgaard N and Mulvany MJ (1994) Number and size of renal glomeruli in Spontaneously Hypertensive Rats. *J. Hypertens.* 12, 1373-1376.
- Swislocki A and Tsuzuki A (1993) Insulin-resistance and hypertension - glucose-intolerance, hyperinsulinemia, and elevated free fatty-acids in the lean Spontaneously Hypertensive Rat. *Am. J. Med. Sci.* 306, 282-286.
- Tanase H (1979) Genetic control of blood pressure in spontaneously hypertensive rats. *Exp. Anim.* 28, 519-530.
- Tanase H and Suzuki Y (1971) Strain difference and genetic determination of blood pressure in rats. *Exp. Animals (Japan)* 20, 1-5.
- Tanase H, Yamori Y, Hansen CT and Lovenberg W (1982) Heart size in inbred strains of rats. Part 1. Genetic determination of the development of cardiovascular enlargement in rats. *Hypertension* 4, 864-872.
- Wang TM, Hsu JF, Jee WSS and Matthews JL (1993) Evidence for reduced cancellous bone mass in the Spontaneously Hypertensive Rat. *Bone and Mineral* 20, 251-264.
- Werner SC, Manger WM, Radichevich I, Wolff M and Von Estorff I (1975) Excessive thyrotropin concentrations in the circulation of the spontaneously hypertensive rat. *Proc. Soc. Exp. Biol. Med.* 148, 1013-1017.
- Wexler BC, McMurtry JP and Iams SG (1981) Histopathologic changes in aging male vs female spontaneously hypertensive rats. *J. Gerontol.* 36, 514-519.
- Womack JE (1973) Biochemical genetics of rat esterases: polymorphisms, tissue expression, and linkage at four loci. *Biochem. Genet.* 9, 13-24.
- Yamori Y, Horie R, Ohtaka M, Nara Y and Ikeda H (1978) Genetic and environmental modification of spontaneous hypertension. *Jpn. Circ. J.* 42, 1151-1159.
- Yamori Y (1984) The development of Spontaneously Hypertensive Rat (SHR) and of various spontaneous rat models, and their implications. In: *Handbook of Hypertension, Vol. 4. Experimental and genetic models of hypertension* (de Jong W, ed). Amsterdam, New York, Oxford: Elsevier, pp 224-239.
- Yamori Y, Nara Y, Tsubouchi T, Sogawa Y, Ikeda K and Horie R (1986) Dietary prevention of stroke and its mechanism in stroke-prone spontaneously hypertensive rats - Preventive effect of dietary fibre and palmitoleic acid. *J. Hypertens.* 4, S449-S452.
- Yamori Y (1994) Development of the spontaneously hypertensive rat (SHR), the stroke-prone SHR (SHRSP) and their various substrain models for hypertension-related cardiovascular disease. In: *Handbook of Hypertension Vol. 16 Experimental and Genetic Models of Hypertension*. (Ganten D and de Jong W, eds). Amsterdam, New York, Oxford: Elsevier, pp. 346-364.
- Yamori Y, Murakami S, Nara Y and Ikeda K (1995) Stroke-prone SHR and arterio-lipidosis-prone SHR as models for atherosclerosis: their mechanisms and application for nutritional and pharmacological studies. *Clin. Exp. Pharmacol. Physiol.* 22, S244-245.
- Yen TT, Roeder H and Willard PW (1974) A genetic study of hypertension in Okamoto-Aoki spontaneously hypertensive rats. *Heredity* 33, 309-316.
- Young JB, Mullen D and Landsberg L (1978) Caloric restriction lowers blood pressure in the spontaneously hypertensive rat. *Metabolism* 27, 1711-1714.