

# SAMP8/TaHsd (Senescence Accelerated Mouse Prone 8)

## SAMP8/TaHsd

The SAMP8 Mouse is developed by Dr. Toshio Takeda, Kyoto University, from AKR/J mice and crossed with mice from an unknown strain, followed by sib mating since 1975.

In 2002 from Takeda Chemical Ltd to Harlan Laboratories. Harlan became Envigo in 2015, then Envigo was acquired by Inotiv in 2021.

## CHARACTERISTICS

### Behavior

SAMP8 mice showed impairments in passive avoidance tasks (Miyamoto *et al*, 1986; Yagi *et al*, 1988), one-way (Miyamoto *et al*, 1986), T-maze (Flood and Morley, 1993) and Sidman (Ohta *et al*, 1989) active avoidance tasks and an impairment of their spatial memory task ability (Miyamoto, 1997). SAMP8 mice had age-related emotional disorders characterized by reduced anxiety-like behavior (Miyamoto *et al*, 1992). The SAMP8 and SAMP10 showed a profound disorder of their circadian rhythms of spontaneous motor activity and drinking behaviors (Miyamoto, 1997).

### Immunology

SAMP8 mice also showed a more rapid decline in the lymphoproliferative response to concanavalin A, but cytotoxic T lymphocyte responses did not change with age. Natural killer cell activity was reduced with age in both SAMR1 and SAMP8, but no strain difference was observed (Powers *et al*, 1995).

### Genetics

#### Coat colour gene

- c : albino.

All SAMP mice have been identified at approximately 20 loci. There is only minor variation among the strains, restricted to *Idh-1*, *Mod-1* and *Car-2* loci.

### Life-span and spontaneous disease

The SAMP8 mouse showed accelerated ageing (Takeda *et al*, 1991). They had age-related emotional disorders characterized by reduced anxiety-like behavior (Miyamoto *et al*, 1992). SAMP8 showed a profound disorder of the circadian rhythm of spontaneous motor activity and drinking behavior (Miyamoto, 1997). The SAMP8 showed age-related appearances of spongiform degeneration in the brain stem (Yagi *et al*, 1989), and of PAS-positive granular structures in their hippocampal formation (Akiyama *et al*, 1986), and astrogliosis in their brain stem (Yagi *et al*, 1989), hippocampus, pyriform cortex, brain stem nuclei and white matter (Kawamata *et al*, 1997). Clusters of activated microglia were also seen around the vacuoles in the brain

stem (Amano *et al*, 1995). A monoamine-oxidase-B-positive granular structure was found in hippocampus of old mice (Nakamura *et al*, 1995). Beta/A4 protein-like immunoreactive granular structures were observed in various regions, including the medial septum, cerebral cortex, hippocampus, cerebellum, and some cranial nerve nuclei and roots and increased markedly in number with age (Takemura *et al*, 1993).

Other age-dependent histological changes included cortical atrophy in the pyriform cortex, neuronal cell loss in the locus ceruleus and lateral tegmental nucleus, intraneuronal accumulation of lipopigment in Purkinje cells, and eosinophilic inclusion bodies in thalamic neurons (Kawamata *et al*, 1997; Akiguchi *et al*, 1994). Similar changes were also observed to a lesser degree in SAMP10.

Blood-brain barrier function was impaired with advancing age in the olfactory bulb and medial hippocampus in SAMP8 (Ueno *et al*, 1993; Ueno *et al*, 1996; Ueno *et al*, 1997; Vorbrodt *et al*, 1995).

SAMP8 mice have a median survival time of 12.1 months.

### Miscellaneous

Characteristics of the SAMP8 strain have been described by Festing (1997) and Lyon *et al*, (1996).





### Physiology and biochemistry

Hippocampal glutamic acid content is higher than in SAMP1 (Nomura *et al*, 1991). Muscarinic acetylcholine receptors, alpha 2-adrenoreceptors, N-methyl-D-aspartate (NMDA) receptor channels and L-type Ca<sup>2+</sup> channels were all changed in the cerebral cortex and hippocampus in aged SAMP8 (Kitamura *et al*, 1989). High levels of K<sup>+</sup> - and NMDA induced L-[<sup>3</sup>H] noradrenalin release in brain slices, and this release was significantly lower in SAMP8 than in SAMR1 (Zhao and Nomura, 1990). Damage to the central histaminergic neurons (Meguro

*et al*, 1992), synaptic dysfunction in the glutamatergic (Kitamura *et al*, 1992) and cholinergic systems (Ikegami *et al*, 1992; Zhao *et al*, 1992) seem to be present in SAMP8. The SAMP8 retained a higher concentration of GM3 than SAMR1 throughout their life span (Ohsawa and Shumiya, 1991). The endogenous levels of the beta-subunit of NGF in these mice was already elevated in the thymus, adrenal gland, testes and hypophysis during the early period of life as compared to SAMR1 (Kato-Semba *et al*, 1991; Kato-Semba *et al*, 1993). Neurotrophin-3 (NT-3) mRNA in the cortex was higher

in SAMP8 than in SAMR1, whereas in the midbrain, hippocampus and forebrain, NT-3 expression levels were lower in SAMP8 than in SAMR1 during early development (Kaisho *et al*, 1994). Brain glucose metabolism was also impaired, as indicated by reductions in 2-deoxyglucose uptake (Kurokawa *et al*, 1996; Ohta *et al*, 1996) and in hexokinase activity (Kurokawa *et al*, 1996) in aged SAMP8. The binding of [<sup>3</sup>H]phorbol-12,13-dibutyrate to protein kinase C in both cytosol and membrane fractions in the hippocampus of aged SAMP8 was reduced (Nomura *et al*, 1997).



## REFERENCES

1. Akiguchi I, Yagi H, Ueno M, Takemura M, Kitabayashi T, Nakamura S, Shimada A, Takeda T (1994) Age-related morphological changes in the brain of Senescence-Accelerated Mouse (SAMP8). In: *The SAM Model of Senescence*. (Takeda T ed). Amsterdam: Excerpta Medica, pp 67-72.
2. Akiyama H, Kameyama M, Akiguchi I, Sugiyama H, Kawamata T, Fukuyama H, Kimura H, Matsuchita M, Takeda T (1986) Periodic Acid-Schiff (PAS)-positive granular structures increase in the brain of senescence accelerated mouse (SAM) *Acta Neuropathol.* 72, 124-129.
3. Amano T, Nakanishi H, Oka M, Yamamoto K (1995) Increased expression of cathepsins E and D in reactive microglial cells associated with spongiform degeneration in the brain stem of Senescence-Accelerated Mouse. *Exp. Neurol.* 136, 171-182.
4. Festing MFW (1997) Inbred Strains of mice. *Mouse genome* 95, 519-686.
5. Flood JE, Morley JE (1993) Age-related changes in foot shock avoidance acquisition and retention in senescence accelerated mouse (SAM). *Neurobiol. Aging* 14, 153-157.
6. Ikegami S, et al (1992) Age-related changes in radial-arm maze learning and basal forebrain cholinergic systems in senescence accelerated mouse (SAM). *Behav. Brain Res.* 51, 15-22.
7. Kaisho Y, Miyamoto M, Shiho O, Onoue H, Kitamura Y, Nomura S (1994) Expression of neurotrophin genes in the brain of senescence-accelerated mouse (SAM) during postnatal development. *Brain Res.* 647, 139-144.
8. Katoh-Semba R, et al (1991) Elevated concentrations of b-nerve growth factor in selected tissues from Senescence Accelerated Mice (SAM-P/8). *Mech. Ageing Dev.* 59, 163-175.
9. Katoh-Semba R, et al (1993) An acceleration of age-related increases in levels of the beta-subunit of nerve growth factor in selected tissues from Senescence-Accelerated Mice (SAM-P/8). *J. Mol. Neurosci.* 4, 107-115.
10. Kawamata T, Akiguchi I, Yagi H, Irino M, Sugiyama H, Akiyama H, Shimada A, Takemura M, Ueno M, Kitabayashi T, Ohnishi K, Seriu N, Higuchi K, Hosokawa M, Takeda T (1997) Neuropathological studies on strains of Senescence-Accelerated Mice (SAM) with age-related deficits in learning and memory. *Exp. Gerontol.* 32, 161-170.
11. Kitamura Y, Zhao XH, Ohnuki T, Nomura Y (1989) Ligand-binding characteristics of [<sup>3</sup>H] QNB, [<sup>3</sup>H] Prazosin, [<sup>3</sup>H] rauwolscine, [<sup>3</sup>H] TCP and [<sup>3</sup>H] nitrendipine to cerebral cortical and hippocampal membranes of senescence accelerated mouse. *Neuroscience Letter* 106, 334-338.
12. Kitamura Y, Zhao XH, Ohnuki T, Takei M, Nomura Y (1992) Age-related changes in transmitter glutamate and NMDA receptor/channels in the brain of senescence-accelerated mouse. *Neuroscience Letter* 137, 169-162.
13. Kurokawa T, Sato E, Inoue A, Ishibashi S (1996) Evidence that glucose metabolism is decreased in the cerebrum of aged female senescence-accelerated mouse, possible involvement of a low hexokinase activity. *Neuroscience Letter* 214, 45-48.
14. Lyon MF, Rastan S, Brown SDM (1996) Genetic variants and strains of the laboratory mouse. 2 Volumes. Oxford, New York, Tokyo: Oxford University Press.
15. Meguro K, Yanai K, Yokoyama K, Sakurai A, Maeyama K, Watanabe T, Matsuzawa T (1992) Neurochemical studies on central histaminergic neuron system of senescence accelerated mouse. *Biogenic Amines* 8, 299-307.
16. Miyamoto M (1997) Characteristics of age-related behavioral changes in Senescence-Accelerated Mouse SAMP8 and SAMPI0. *Exp. Gerontol.* 32, 139-148.
17. Miyamoto M, Kiyota Y, Nishiyama M, Nagaoka A (1992) Senescence-accelerated mouse (SAM): age-related reduced anxiety-like behavior in the SAM-P/8 strain. *Physiol. Behav.* 51, 979-985.
18. Miyamoto M, Kiyota Y, Tamazaki N, Nagonka A, Matsuo T, Noga Y, Takeda T (1986) Age-related changes in learning and memory in the Senescence Accelerated Mouse (SAM). *Physiol. Behav.* 45, 393-406.
19. Nakamura S, Akiguchi I, Seriu N, Ohnishi K, Takemura M, Ueno M, Tomimoto H, Kawamata T, Kimura J, Hosokawa M (1995) Monoamine oxidase-B-positive granular structures in the hippocampus of aged senescence-accelerated mouse (SAMP8). *Acta Neuropathol.* 90, 626-632.
20. Nomura Y, Kitamura Y, Ohnuki T, Arima T, Yamanaka Y, Sasaki K, Oomura Y (1997) Alterations in acetylcholine, NMDA, benzodiazepine receptors and protein kinase C in the brain of senescence-accelerated mouse: an animal model useful for studies on cognitive enhancers. *Behav. Brain Res.* 83, 51-55.
21. Nomura Y, Kitamura Y, Zhao XH (1991) Aging in glutamergic system with special reference the NMDA receptor/ion channel complex in the brains of senescence accelerated mice. In: *NMDA Receptor related agents: Biochemistry, Pharmacology and Behavior* (Kameyama T, Domin F, Nabeshima, eds) Ann Arbor, MI: NPP books, pp 287-298.
22. Ohsawa T, Shumiya S (1991) Age-related alteration of brain gangliosides in Senescence-Accelerated Mouse (SAM)-P/8. *Mech. Ageing Dev.* 59, 263-274.
23. Ohta A, Hirano T, Yagi H, Tanaka S, Hosokawa M, Taneda T (1989) Behavioral characteristics of the SAM-P/8 strain in Sidman active avoidance Task. *Brain Res.* 498, 195-198.
24. Ohta H, Nishikawa H, Hirai K, Kato K, Miyamoto M (1996) Relationship of impaired brain glucose metabolism to learning deficit in senescence-accelerated mouse. *Neurosci. Letter* 217, 37-40.
25. Powers DC, et al (1995) Age-related changes in lymphoproliferation, cytotoxic T lymphocyte responses, and natural killer activity of Senescence-Accelerated Mouse (SAM)-P/8 and SAM-R/1 substrains. *Aging. Immunol. Infec. Dis.* 6, 43-52.
26. Takeda T, Hosokawa M, Higuchi K (1991) Senescence-accelerated mouse (SAM): A novel murine model of accelerated senescence. *J. Amer. Geriatr. Soc.* 39, 911-919.
27. Takemura M, Nakamura S, Akiguchi I, Ueno M, Oka N, Ishikawa S, Shimada A, Kimura J, Takeda T (1993) Beta/A4 protein like immunoreactive granular structures in the brain of Senescence-Accelerated Mouse (SAM). *Am. J. Pathol.* 142, 1887-1897.
28. Ueno M, Akiguchi I, Hosokawa M, Shinnou M, Sakamoto H, Takemura M, Higuchi K (1997) Age-related changes in the brain transfer of blood-borne horseradish peroxidase in the hippocampus of Senescence-Accelerated Mouse (SAM). *Acta Neuropathol.* 93, 233-240.
29. Ueno M, Dobrogowska DH, Vorbrodt AW (1996) Immunocytochemical evaluation of the blood-brain barrier to endogenous albumin in the olfactory bulb and pons of senescence accelerated mice (SAM). *Histochem. Cell Biol.* 105, 203-212.
30. Ueno M, et al (1993) Age-related changes in barrier function in mouse brain. I. Accelerated age-related increase of brain transfer of serum albumin in accelerated senescence prone SAM-P/8 mice with deficits in learning and memory. *Arch. Gerontol. Geriatr.* 16, 233-248.
31. Vorbrodt AW, et al (1995) A quantitative immunocytochemical study of blood-brain barrier to endogenous albumin in cerebral cortex and hippocampus of senescence-accelerated mice (SAM). *Folia Histochem. et Cytobiol.* 33, 229-237.
32. Yagi H, Irino M, Matsushita T, Katoh S, Urmezawa M, Tsuboyama T, Hosokawa M, Akiguchi I, Tokunaga R, Takeda T (1989) Spontaneous spongy degeneration of the brain stem in SAM-P/8 mice, a newly developed memory-deficient strain. *J. Neuropathol. Exp. Neurol.* 48, 577-590.
33. Yagi H, Katoh S, Akiguchi I, Taneda T (1988) Age-related deterioration of ability of acquisition in memory and learning in senescence accelerated mouse: SAM-P/8 as an animal model of disturbance in recent memory. *Brain Res.* 474, 86-93.
34. Zhao X-H, Kitamura Y, Nomura Y (1992) Age-related changes in NMDA-induced [<sup>3</sup>H] acetylcholine release from brain slices of Senescence-Accelerated Mouse. *Int. J. Dev. Neuroscience* 10, 121-129.
35. Zhao X-H, Nomura Y (1990) Age-related changes in uptake and release on L-[<sup>3</sup>H] noradrenaline in brain slices of Senescence Accelerated Mouse. *Int. J. Dev. Neuroscience* 8, 267-272.